Land use and cover change scenarios in the Mesoamerican Biological Corridor-Chiapas, México

DIANA RAMÍREZ-MEJÍA, GABRIELA CUEVAS, PAULA MELI, EDUARDO MENDOZA

Supplementary Information

Table 1s. Extent and rates of change of the different land use and cover categories in the Mesoamerican Biological Corridor-Chiapas in the different periods evaluated.

Land cover class	1993		2002		2007		LUCC rates	
	ha	%	ha	%	ha	%	1993-2002 (%/year)	2002-2007 (%/year)
TroF	837525	23.29	662500	18.42	626125	17.41	-2.57	-0.80
TroSF	707500	19.68	772850	21.49	741550	20.62	0.99	-0.59
TemF	626425	17.42	421800	11.73	360200	10.02	-4.30	-2.23
TemSF	355225	9.88	520850	14.48	560125	15.58	4.34	1.04
ALF	1003550	27.91	1151475	32.02	1243750	34.59	1.54	1.11
OC	65675	1.83	66425	1.85	64150	1.78	0.13	-0.50
Total	3595900	100	3595900	100	3595900	100

Figure 1s. Weights of evidence calculated for the variables: a) protected area presence and b) distance to roads, for specific transitions. TroF = Closed tropical forests, TroSF = Secondary tropical forests, TemF = Closed temperate forests, TemSF = Secondary temperate forests, ALF = Agriculture, livestock and forestry, OC = Other land cover.

Figure 2s. Fuzzy similarity index between the real and simulated maps as a function of spatial resolution.