SEED TRAITS AND GERMINATION IN THE CACTACEAE FAMILY: A REVIEW ACROSS THE AMERICAS

RASGOS SEMINALES Y GERMINACIÓN EN LA FAMILIA CACTACEAE: UNA REVISIÓN EN LAS AMÉRICAS

DUNIEL BARRIOS^{1*}, JORGE A. SÁNCHEZ², JOEL FLORES³, ENRIQUE JURADO⁴

¹Jardín Botánico Nacional, Universidad de La Habana, Cuba.

²Instituto de Ecología y Sistemática, Ministerio de Ciencia, Tecnología y Medio Ambiente, Cuba.

³División de Ciencias Ambientales, Instituto Potosino de Investigación Científica y Tecnológica, AC. México.

⁴Facultad de Ciencias Forestales, Universidad Autónoma de Nuevo León, Linares, México

*Corresponding author: duniel.barrios@gmail.com

Abstract

Background: Cactaceae is the fifth taxonomic group with the highest proportion of threatened species. One way to contribute to the preservation of this family is to understand the processes that promote seed germination.

Questions: How common is dormancy and seed banks in Cactaceae? Are there general patterns in cacti germination response to temperature, light, water, salinity, phytohormones, hydration/dehydration cycles, mechanical or chemical scarification?

Data description: A total of 333 studies on cactus germination with information on 409 taxa.

Study site and dates: since 1939 to January 2020.

Methods: A search of scientific articles in Google Scholar was performed with the words Cactaceae, cacti and cactus, in combination with various matters on germination in English, Spanish and Portuguese.

Results: The main germination studies in cactus deal with photoblasticism (275 taxa), temperature (205 taxa) and seed longevity (142 taxa). Other lines of study in cactus germination (*e.g.*, desiccation tolerance, vivipary, phytohormones, mechanical or chemical scarification, *in vitro* germination, hydration/dehydration cycles, water and saline stress, serotiny, storage in cold, high temperature tolerance and soil seed bank) include between 14 and 65 taxa. Cacti have only physiological dormancy and optimal germination for most species occur between 20 and 30 °C.

Conclusions: Mexico, Brazil and Argentina are the three leading countries in the study of cactus germination.

Keywords: Dormancy, photoblasticism, seed bank, serotiny, temperature, viviparity.

Resumen:

Antecedentes: Cactaceae constituye el quinto grupo taxonómico con mayor proporción de especies amenazadas. Uno de los aspectos que contribuyen a la preservación de estas especies es el entendimiento de los procesos que promueven la germinación de sus semillas.

Preguntas: ¿Qué tan común es la latencia y los bancos de semillas en Cactaceae? ¿Existen patrones generales en la respuesta germinativa de los cactus ante la temperatura, luz, agua, salinidad, fitohormonas, ciclos de hidratación/deshidratación o escarificación mecánica o química?

Descripción de datos: Se revisaron 333 estudios sobre germinación de cactus con información de 409 taxa.

Sitio y años de estudio: Estudios de germinación de cactáceas en América, publicados desde 1939 a enero de 2020.

Métodos: Se realizó una búsqueda de artículos científicos en Google Académico con las palabras Cactaceae y cactus, en combinación con varias materias sobre germinación en inglés, español y portugués.

Resultados: Los principales estudios sobre la germinación de cactáceas han versado sobre el fotoblastismo (275 taxones), la temperatura (205 taxones) y la longevidad seminal (142 taxones). Otras líneas de estudio que abordan la germinación de cactus (*e.g.*, tolerancia a la desecación, viviparidad, fitohormonas, escarificación mecánica o química, germinación *in vitro*, ciclos de hidratación/deshidratación, estrés hídrico y salino, serotinia, tolerancia a altas temperaturas y almacenamiento en frío, bancos de semillas) comprenden entre 14 y 65 taxones. Las cactáceas presentan solo latencia fisiológica y una temperatura óptima de germinación entre los 20 y 30 °C.

Conclusiones: México, Brasil y Argentina son los tres países líderes en el estudio de la germinación en cactus.

Palabras claves: Banco de semillas, fotoblastismo, latencia, serotinia, temperatura, viviparidad.

Cactaceae is the family considered to have the largest number of genera within the order Caryophyllales (Hernández-Ledesma *et al.* 2015) and its species occupy a wide range of habitats through the American continent (Anderson 2001, Goettsch *et al.* 2015). For centuries cacti have fascinated humans because of the beauty and the rarity of the majority of their species (Bravo-Hollis 1978, Griffith 2004). Although the most widespread use is ornamental (Goettsch *et al.* 2015), in several countries the fruits, flowers and stems of 154 species are widely consumed (Anderson 2001, Goettsch *et al.* 2015). Plants of the Cactaceae are currently among the most vulnerable to human disturbance because of their slow growth (Godínez-Álvarez *et al.* 2003) and constitute the fifth taxonomic group with the highest proportion of species threatened worldwide (Goettsch *et al.* 2015).

One way to help preserve these species and others with special conservation value is to understand the processes that promote seed germination (Rojas-Aréchiga & Vázquez-Yanes 2000, Anderson 2001, Flores *et al.* 2006). Two remarkable works about cactus seeds have been published. Rojas-Aréchiga & Vázquez-Yanes (2000) published an exhaustive review on cactus seed germination. Barthlott & Hunt (2000) published a book on cactus seed micromorphology, including 213 species and 688 microphotographs from more than 350 taxa. In addition, it contains the compilation of studies with microphotographs of cactus seeds from 1,050 species of 230 genera (Rojas-Aréchiga 2012). Barthlott & Hunt (2000) study describes 26 useful seed traits and attempts to homogenize English terminology. Both Rojas-Aréchiga & Vázquez-Yanes (2000) and Barthlott & Hunt (2000) studies have been referenced for all research in cactus seed germination and seed description during the last two decades.

Since the 2000s, the papers on cactus seed germination almost tripled the available information. This review aims to perform an analysis of the cactus seed literature in the last 80 years. The trends found and gaps in the information are discussed. Specifically, we analyzed the relationship between cactus seed germination and seed traits, as well as between cactus seed germination and phylogeny. We also documented the germination types in cacti, and how common is seed dormancy and how common are seed banks in the cactus family. We also discuss the effect of temperatures, light, water, salt and hydration-dehydration cycles on cactus seed germination.

Materials and methods

A search of scientific articles was performed with the words Cactaceae, cacti or cactus in combination with: germination, photoblasticism, seed bank, vivipary,

dormancy, hydration-dehydration cycle, serotiny and *in vitro* germination through the search engine Google Scholar (<https://scholar.google.com>), in English, Spanish, and Portuguese, the main languages in the American continent. According to Haddaway *et al.* (2015) Google Scholar is a powerful tool for finding specific literature and is adequate for identifying the majority of evidence in a systematic review, the references from each study found were also analyzed. In total 333 studies related to cactus germination from 1939 to January 2020 (Figure 1, Tables 1 and 2) were gathered, with information on 409 taxa. Cactus seed germination has been evaluated in less than 50 % of the genera. The collected information was grouped by subjects and then discussed.

Figure 1. Number of studies of cactus seed germination until January, 2020.

All the information obtained was divided into ten subjects according topics covered in Baskin & Baskin (2014) and that there were also several studies on cacti. In the section on *Cactus seed traits*; shape, color, size, mass, as well as appendices and number of seeds per fruit were considered. The intention of this topic was to introduce readers to the generalities of cactus seeds as a necessary process to discuss the different issues related to germination. Also the section on *Germination types*, that relates the shape of the seed to the mode of germination, was mainly considered as introductory. The information discussed in these two subjects was obtained mainly from classical or compilation studies (e.g., Flores & Engelman 1976, Bregman & Bouman 1983, Barthlott & Hunt 2000, Rojas-Aréchiga & Vázquez-Yanes 2000, Seal *et al.* 2009).

The number of taxa studied by genera, subfamilies and country or region of origin of the taxon was compiled for the rest of the subjects. *Seed dormancy*, was not included in the summary table because we used the number of cactus species having seed dormancy from Willis *et al.* (2014). The species where viviparity has been studied were also included in the summary table.

Table 1. Abstract of the cactus studies on different research lines related with seed germination in 80 years and 333 papers with information on 409 taxa. Taxa: Number of taxa studied; Cact: *Cactoideae*; Opt: *Opuntioideae*; Per: *Pereskioideae*; Mah: *Maihuenioideae*. Countries or regions, MX: Mexico; BR: Brazil; AR: Argentina; CH: Chile; PE: Peru; BO: Bolivia; PA: Paraguay; CO: Colombia; USA: United States of America; CA: Caribbean region; SOU: South America; MX-U: Mexico - USA; CU: Cuba. It is possible to find a higher number of species per country when in a same taxon several studies in natural populations from different countries were done. In parenthesis the percentage of genera with at least one species studied.

Matter	Genus	Taxa	Subfamilies					Countries or regions											
			Cact	Opt	Per	Mah	MX	BR	AR	CH	PE	BO	PA	CO	USA	CA	SOU*	MX-U*	CU*
Photoblasticism	62 (50.0)	275	251	20	3	1	129	46	41	37	9	0	0	0	7	4	0	0	16
Temperatures	62 (50.0)	205	184	17	3	1	64	43	30	14	12	4	1	0	14	10	13	0	17
Longevity ¹	43 (34.7)	142	128	12	1	1	70	25	10	19	2	1	0	0	11	2	7	0	9
Desecation tolerance	30 (24.2)	65	53	2	0	0	19	0	6	13	6	1	0	0	1	2	10	7	0
Vivipary	21 (16.9)	58	58	0	0	0	12	5	27	0	0	1	0	0	1	6	6	0	13
Phytohormones (AG ₃)	29 (23.4)	59	50	9	0	0	36	4	4	4	0	0	1	0	6	2	4	0	10
Mechanical or chemical scarification ²	23 (18.5)	52	41	11	0	0	40	2	2	0	0	0	0	0	7	0	1	0	12
<i>In vitro</i> germination	18 (14.5)	42	38	4	0	0	26	14	0	0	0	0	0	0	0	3	0	0	0
Water stress	23 (18.5)	39	37	1	1	0	15	10	9	2	0	0	0	3	0	0	0	0	9
Serotiny	9 (7.2)	24	22	2	0	0	20	0	0	0	0	0	0	0	3	1	0	0	1
Hydration/dehydration cycles	11 (8.8)	20	20	0	0	0	11	2	7	0	0	0	0	0	0	0	0	0	1
Storage in cold ³	7 (5.6)	20	20	0	0	0	2	16	0	0	0	0	0	0	2	0	0	0	0
Soil seed bank	10 (8.1)	19	17	2	0	0	7	3	6	0	0	0	0	0	3	0	0	0	9
High temperature tolerance	9 (7.2)	17	13	4	0	0	14	1	0	0	0	0	0	0	3	0	0	0	0
Saline stress	8 (6.4)	14	12	2	0	0	7	7	0	0	0	0	0	0	0	0	0	0	8

¹ It includes studies where the viability of seeds of at least three months stored at room temperature was tested.

² It includes studies where the germination response was tested after subjecting the seeds to various scarification methods that include sand, sandpaper, cuts (mechanical scarification) or with H₂SO₄, HCl, NaClO and H₂O₂ (chemical scarification).

³ Only studies where the seeds were kept at temperatures equal to or below 0 °C were included.

* The species referred for SOU and MX-U comprise species where the seed collection site was not declared and the distribution covers two or more countries. The references for Cuba were not added to the total species, because they include studies that have not yet been published, the germination studies of Cuban cacti that have already been published were included in the Caribbean list.

In the subject *Temperature and seed germination*, the mean and the confidence intervals of the germination percentage at 30 °C were plotted between cacti from temperate and warm climates. The type of weather was obtained from the information of each article reviewed; the comparison was based on 43 data from temperate and 34 from warm climate. Similarly, in the subject *Water and saline stress and germination*, the mean and standard deviation of the germination response of cactus seeds were plotted against different water potentials induced with polyethylene glycol (PEG).

On the subject *Light and seed germination*, we consider the photoblastic response of the seeds according to the Relative Light Germination index (RLG) (Milberg *et al.* 2000) and the proportions proposed by Funes *et al.* (2009).

Values with RLG > 0.75 were considered positive photoblastic, < 0.25 negative photoblastic, species with RLG values between 0.25 and 0.75 were considered indifferent photoblastic (Funes *et al.* 2009). We calculate the RLG, to define the photoblastic response of the seeds in the studies where they had not calculated it.

Results and Discussion

Cactus seed traits. Both the works of Flores & Engelman (1976) and Bregman & Bouman (1983), as well as of Barthlott & Hunt (2000), present the structure and general anatomy of cacti seeds. In such studies, it was shown that in cacti the embryo is covered by two layers of tissue: the external (known as the testa) is generally thick, while the

Seed traits and germination in the Cactaceae

Table 2. Taxa of the *Cactaceae* having at least nine species with seed germination studies. Taxonomy was based in [Hunt et al. \(2006\)](#). Spp = Species number per genus. Review based in 333 papers published in the last 80 years.

Subfamily/genus	spp	With information	Subfamily/Genus	spp	With information
<i>Cactoideae</i>			<i>Cactoideae</i>		
<i>Armatocereus</i>	9	1	<i>Melocactus</i>	50	18
<i>Browningia</i>	9	0	<i>Micranthocereus</i>	10	2
<i>Cereus</i>	29	6	<i>Pachycereus</i>	13	6
<i>Cleistocactus</i>	48	5	<i>Parodia</i>	66	8
<i>Coleocephalocereus</i>	10	1	<i>Pediocactus</i>	10	0
<i>Copiapoa</i>	30	6	<i>Peniocereus</i>	20	2
<i>Corryocactus</i>	12	1	<i>Pfeiffera</i>	10	0
<i>Coryphantha</i>	53	9	<i>Pilosocereus</i>	49	18
<i>Discocactus</i>	12	3	<i>Pseudorhipsalis</i>	9	0
<i>Disocactus</i>	16	1	<i>Rebutia</i>	40	1
<i>Echinocereus</i>	106	4	<i>Rhipsalis</i>	48	9
<i>Echinopsis</i>	101	28	<i>Sclerocactus</i>	28	2
<i>Epiphyllum</i>	18	4	<i>Selenicereus</i>	27	1
<i>Eriosyce</i>	51	17	<i>Stenocereus</i>	25	9
<i>Escobaria</i>	23	5	<i>Thelocactus</i>	20	3
<i>Espostoa</i>	12	2	<i>Turbinicarpus</i>	36	22
<i>Ferocactus</i>	42	13	<i>Weberocereus</i>	10	1
<i>Frailea</i>	18	1	<i>Opuntioideae</i>		
<i>Gymnocalycium</i>	63	9	<i>Austrocylindropuntia</i>	9	0
<i>Haageocereus</i>	14	5	<i>Corynopuntia</i>	14	0
<i>Harrisia</i>	9	7	<i>Cylindropuntia</i>	33	5
<i>Hylocereus</i>	14	7	<i>Opuntia</i>	75	17
<i>Leptocereus</i>	11	1	<i>Pterocactus</i>	9	1
<i>Mammillaria</i>	232	43	<i>Pereskioideae</i>		
<i>Matucana</i>	21	0	<i>Pereskia</i>	18	3

internal one only forms a fine membrane around the embryo ([Bregman & Bouman 1983](#)) that completely collapses during seed maturation in most species ([Barthlott & Hunt 2000](#)) ([Figure 2](#)). A distinctive trait in *Cactaceae* seeds, but not exclusive of the family, is a small area called the hilum-micropylar region, which is caused by the presence of campylotropous ovules ([Bregman & Bouman 1983](#), [Barthlott & Hunt 1993](#)), although [Flores & Engleman \(1976\)](#) considered the amphitropous ovules dominant. This characteristic is considered a variant of campylotropous ovules according to [Font Quer \(1973\)](#). For descriptive purposes [Barthlott & Hunt \(2000\)](#) divided cactus seeds in four regions: ventral, dorsal, apical and the edge of the hilum-micropylar region.

Cactaceae seeds vary in shape. [Bregman & Bouman \(1983\)](#) in their review presented 12 seed variants, while [Barthlott & Hunt \(2000\)](#) in accordance with the treatment of [Voit \(1979\)](#) considered eight types of seeds, which they

themselves reduced to four, according to the length/width relationship. In general, cactus seeds have been described as reniform, pyriform, globular, oval, hat-shaped, lenticular or mussel-shaped ([Rojas-Aréchiga & Vázquez-Yanes 2000](#), [Barthlott & Hunt 2000](#)) ([Figure 3](#)). In spite of the great variability of terminologies, the majority of the previous studies have followed [Barthlott & Hunt \(2000\)](#) (see: [Arias & Terrazas 2004](#), [Arroyo-Consultchi et al. 2006, 2007](#), [Arias et al. 2012](#), and [Franco-Estrada et al. 2014](#)). Although it is possible that some relationship between the shape of the cactus seeds and their germination exists, this aspect has been little studied. [Bregman & Bouman \(1983\)](#) associated the seed shape with differences in the formation of germination cracks. Hat-shaped seeds occur in North American genera such as *Astrophytum* and *Epithelanta*, as well in species of several South American genera, such as *Frailea*, *Matucana*, *Discocactus*, *Notocactus* (*Parodia*), and *Thrixanthocereus* (*Espostoa*) ([Barthlott & Hunt 2000](#)).

Experimental evidence of floating capabilities (hat shape, a funicular envelope covering a prominent hilum, and air chambers throughout the tegument) and potential water dispersal of hat-shaped seeds in Cactaceae have been found in *Astrophytum capricorne* and *Astrophytum ornatum* (Sánchez-Salas *et al.* 2012), as well as in *Astrophytum myriostigma* (Romero-Méndez *et al.* 2018). Seeds of *Selenicereus wittii* from the Amazonia, in Brasil, are mussel-shaped, and also have an air chamber and are adapted to inundation hydrochory (Barhlott *et al.* 1997).

Figure 2. Transversal section of a cactus seed. t: testa; il: internal layer and e: embryo. Drawing by Alfredo Ruiz Fleitas

Unlike the high variability in terms of shape, seed color in the family is mainly between black and brown (Leuenberger 1986, Barhlott & Hunt 2000, Arias & Terrazas 2004, Arroyo-Consultchi *et al.* 2006, Seal *et al.* 2009, Arias *et al.* 2012, Loza-Cornejo *et al.* 2012, Franco-Estrada *et al.* 2014). In some cases; however, they can be reddish-brown, yellow, green or off-white (Rojas-Aréchiga & Vázquez-Yanes 2000, Barhlott & Hunt 2000, Arias *et al.* 2012). A potential relationship between seed color and seed germination in cactus remains to be tested.

Consistent with seed shape, the embryo of the cactus is generally curved (Barhlott & Hunt 2000) and peripheral (Finch-Savage & Leubner-Metzger 2006), with great variation in the region that occupies the storage structure. In *Leuenbergeria*, *Opuntia* and *Pereskopsis* species, a greater development of the cotyledons is observed with respect to the hypocotyl. Nevertheless, the tendency in the family is a greater development of the hypocotyl with respect to the cotyledons. Cactus genera like *Mammillaria* and *Parodia* show upright embryos with 90 % of the volume occupied by the hypocotyl (Barhlott & Hunt 2000).

In terms of the average number of seeds produced per fruit, an enormous variation exists in cacti. Some authors report cactus fruits (e.g. *Epiphyllum anguliger*, *Ferocactus*

histris, *Pachycereus pringlei*, *P. weberi*, *Pilosocereus chrysacanthus*, *Echinopsis atacamensis* subsp. *pasacana*, *Gymnocalycium monvillei*), with ranges between 1,000 and 8,000 seeds (Zimmer 1966, Del Castillo 1988, Fleming *et al.* 1994, Valiente-Banuet *et al.* 1997, de Viana 1999, Gurvich *et al.* 2008), although it is possible to find species with fewer than 20 seeds per fruit in genera such as *Mammillaria* (Valverde & Zavala-Hurtado 2006, Valverde *et al.* 2015), *Pereskia* and *Leuenbergeria* (Leuenberger 1986).

Cactus seeds are generally smaller than one centimeter (Barhlott & Hunt 2000, Seal *et al.* 2009, Rojas-Aréchiga *et al.* 2013) and vary from 0.2 mm in *Blossfeldia liliputana* (Barhlott & Porembski 1996) to 7.5 mm in *Pereskia bleo* (Leuenberger 1986), and their weight varies from 0.016 mg in *Mammillaria bocasana* (Flores *et al.* 2006) to 56.04 mg in *Opuntia basilaris* (Royal Botanic Gardens Kew 2019).

Differences in seed mass among cactus species and localities have been found, e.g., Romo-Campos *et al.* (2010) found that *Opuntia* spp. seeds collected in moist areas were heavier than those collected in dry sites. These authors also found a correlation between seed mass and seed germination for *Opuntia jaliscana* in which larger seeds germinated more than smaller seeds. Sosa-Pivatto *et al.* (2014) found no correlation between seed mass and germination characteristics in central Argentinian cacti; however, they found that species with heavier seeds produced bigger and more cylindrical seedlings.

Other traits associated with cactus seeds are appendages or structures, such as funicula, arils, strophioles and mucilage. In the subfamily Opuntioideae, the ovule is surrounded by the funiculus that is lignified during development (Bregman & Bouman 1983, Strittmatter *et al.* 2002) and forms a structure considered as an aril by Barhlott & Hunt (2000) but as a covering by Bravo-Hollis (1978). Orozco-Segovia *et al.* (2007) found that lignified funiculus in *Opuntia tomentosa* acts as a partial barrier to water diffusion into the seed and that in this species, water uptake occurs mainly through the water channel mediated by a valve, both formed in the hilum-micropyle region during seed dehydration and ageing. Various genera like *Mammillaria*, *Aztekium* (Bravo-Hollis 1978) and *Blossfeldia* (Barhlott & Hunt 2000) show strophioles that can be considered arils, but to differentiate them from the structure of Opuntioideae, this term has continued to be used (Barhlott & Hunt 2000). A mucilage sheath is present in more than 40 species of cactus belonging to 22 genera and is formed by pectins (Barrios *et al.* 2015, Mascot-Gómez *et al.* 2020). Bregman & Graven (1997) and Mascot-Gómez *et al.* (2020) suggested that the layer of mucilage in cactus seeds improves the germination through the effect of the intake and distribution of water. Barrios *et al.* (2015) suggested that maybe the presence of mucilage in

Figure 3. Shape of cactus seeds. A: globular (*Blossfeldia liliputana*); B: lenticular (*Leuenbergeria bleo*); C: pyriform (*Escobaria cubensis*); D: mussel-shaped (*Selenicereus grandiflorus*); E: reniform (*Neobuxbaumia multiareolata*); F: hat-shaped (*Frailea phaeodisca*). According [Barthlott & Hunt \(2000\)](#) A and B: circular; C and D: oval; E and F: broadly oval. Drawings by Alfredo Ruiz Fleitas

Leptocereus scopulophilus seeds serves other purposes such as the adhesion of seeds to the soil or camouflage from seed eaters.

Germination types. Seed germination is a series of events that begin with imbibition and end with the emergence of the radicle from the seed coat. There are few studies that consider germination types in Cactaceae ([Almeida et al. 2009](#), [Secorun & Souza 2011](#)), since the classic study by [Bregman & Bouman \(1983\)](#). The aforementioned authors described 11 variants of germination where the radicle is the first organ of the seedling to break the testa and hypothesized the phylogenetic relationships between variants. In addition, they considered that the mode of germination in a cactus is correlated to the seed shape, and they defined the *Cereus* variant as the most numerous in the family. Curiously, although for the genus *Leptocereus* [Areces-Mallea \(2003\)](#) only recognized the *Cereus* variant in two species (*L. quadricostatus* and *L. sylvestris*), [García-Beltrán et al. \(2017\)](#) found four variants in only one species (*L. scopulophilus*). In addition to the *Cereus* variant in *L. scopulophilus*, [García-Beltrán et al. \(2017\)](#) recognized the

Pereskia variant and another two where the cotyledons are the first organ to break the testa. The aforementioned variants were considered exceptions by [Bregman & Bouman \(1983\)](#). Nevertheless, in one of the seed morphs of *L. scopulophilus*, these “alternative variants” account for over 20 % ([García-Beltrán et al. 2017](#)).

Seed dormancy. [Rojas-Aréchiga & Vázquez-Yanes \(2000\)](#) only found reports of seeds with innate and forced dormancy. The forced dormancy of seeds is considered by [Baskin & Baskin \(2014\)](#) as non-dormant. The classification proposed by those authors establishes that seeds that germinate within the first four weeks after dispersion are non-dormant and that the seeds that do not germinate because of inadequate conditions (forced dormancy) must be considered as latent or quiescent. Previous cactus studies have reported viviparous seeds (without innate dormancy) in 58 species ([Table 1](#)) ([Cota-Sánchez 2004](#), [Cota-Sánchez et al. 2007, 2011](#), [Rojas-Aréchiga & Mandujano-Sánchez 2009](#), [Barrios et al. 2012](#), [Aragón-Gastélum et al. 2013, 2017](#)). Viviparous are also considered as non-dormant in the system of [Baskin & Baskin \(2014\)](#). The studies about cactus

seed dormancy are few ([Potter *et al.* 1984](#), [Crtes *et al.* 1994](#), [Olvera-Carrillo *et al.* 2003](#), [Mandujano *et al.* 2005](#), [Flores *et al.* 2005](#), [2008](#), [Orozco-Segovia *et al.* 2007](#), [Delgado-Snchez *et al.* 2010](#), [2011](#), [2013](#)), may be because of the rapid germination that the majority of the studied species possess. In addition, few works about germination mention how many un-germinated seeds are dormant ([Flores *et al.* 2005](#), [2006](#), [Ortega-Baes *et al.* 2010](#), [Aragn & Lasso 2018](#)).

The most used methods for interrupting dormancy in cactus seeds have been through chemical scarification (by immersion in H₂SO₄, HCl, NaClO and H₂O₂ at various concentrations and times) and mechanical scarification, which together encompass some 52 species studied ([Table 1](#)). In addition, gibberellic acid has been used as a promoter of germination, but its role is discussed later in reference to light. Few studies have used thermal shock but without obtaining good results with this type of treatment ([Snchez-Venegas 1997](#), [Rosas-Lpez & Collazo-Ortega 2004](#), [Carbajal *et al.* 2010](#), [Villanueva *et al.* 2016](#), [Podda *et al.* 2017](#), [Gonzalez-Corts *et al.* 2018](#)). Other studies have used fungal inoculation in some species of *Opuntia* ([Delgado-Snchez *et al.* 2010](#), [2011](#), [2013](#)) or hydration/dehydration cycles ([Dubrovsky 1996](#), [1998](#), [Santini & Martorell 2013](#), [Contreras-Quiroz *et al.* 2016a](#), [2016b](#)) with good results. Research in the last decade that has addressed treatments to break dormancy in cactus seeds are however, not common ([Rojas-Archiga *et al.* 2011](#), [Amador-Alfrez *et al.* 2013](#), [Delgado-Snchez *et al.* 2010](#), [2011](#), [2013](#), [Podda *et al.* 2017](#), [Gonzalez-Corts *et al.* 2018](#)).

In cacti, a rapid and high percentage of germination is associated with a thin testa ([Maiti *et al.* 1994](#)). However, although testa functions as regulator in imbibition ([Souza & Marcos-Filho 2001](#)), physical dormancy has not been found in cactus seeds. Although works about germination in *Opuntia* and *Harrisia* have suggested the presence of physical, morphological or morphophysiological dormancy ([Potter *et al.* 1984](#), [Olvera-Carrillo *et al.* 2003](#), [Podda *et al.* 2017](#)), morphological ([Mandujano *et al.* 1997](#)) or morphophysiological ([Dehan & Prez 2005](#)). None of these studies demonstrated a lack of seed imbibition, or an undifferentiated or underdeveloped embryo, which are traits used to show the existence of these types of dormancy ([Baskin & Baskin 2014](#)). Previous studies of *Opuntia* (including the species erroneously classified with physical and morphological dormancy) have demonstrated that only physiological dormancy can be attributed to them ([Mandujano *et al.* 2005](#), [Orozco-Segovia *et al.* 2007](#)). In addition, cactus seeds having physiological dormancy can show dormancy cycling, [Aragn-Gastlum *et al.* \(2018\)](#) found that buried seeds of *Echinocactus platyacanthus* acquired secondary dormancy in the rainy seasons (summer

and autumn), which was alleviated at the end of the subsequent dry season (winter), possibly because of the high variation registered in mean and minimum soil temperature at the end of winter.

[Finch-Savage & Leubner-Metzger \(2006\)](#), [Willis *et al.* \(2014\)](#) and [Baskin & Baskin \(2014\)](#) currently only recognize non-dormant seeds (including viviparous) and seeds with physiological dormancy for Cactaceae. According to [Willis *et al.* \(2014\)](#), in 153 cactus species studied 65 % showed physiological dormancy and 35 % non-dormant seeds. Most cacti inhabit arid and semiarid environments, so it is expected that future studies will find new species with physiological dormancy and probably species with physical dormancy as an adaptation to delay of germination, as has been proposed for species that occupy extreme or seasonal habitats ([Jurado & Flores 2005](#), [Baskin & Baskin 2014](#)). In addition, a greater presence of heteromorphism in Cactaceae seeds with different degrees of physiological dormancy could be expected, as it occurs in *Leptocereus scopulophilus* ([Garca-Beltrn *et al.* 2017](#)) and *Leptocereus arboreus* seeds (unpublished data). Seed heteromorphism is considered a “bet hedging” strategy that favors the reproductive success of the offspring in unpredictable and heterogeneous environments ([Venable 2007](#)).

Seed banks. A seed bank is formed through the presence of viable seeds that remain in the habitat without germinating during a given period ([Thompson & Grime 1979](#)). Two types of banks have been defined in terms of the longevity of the seeds: a transient seed bank, when the seeds do not stay viable for more than a year ([Thompson & Grime 1979](#)) or until the next germination season ([Walck *et al.* 2005](#)) and a persistent seed bank when the seeds achieve permanence for more than a year.

When the place where the seeds are stored is referred to, two types of banks have been described: aerial and terrestrial ([Baskin & Baskin 2014](#)). Seeds are typically stored in the soil (the soil or terrestrial seed bank) but, in some species, seeds may be retained in plant canopy (the aerial seed bank) until seed release is triggered ([Gnster 1992](#)). The formation of seed banks requires from the species an orthodox storage behavior that allows them to remain viable for long periods, as well as to escape from predators ([Baskin & Baskin 2014](#)).

The first studies about cactus seed banks concluded their existence whether through the collection of viable seeds in the soil ([de Viana 1999](#)), by examining seeds viability after different burial periods ([Bowers 2000](#), [2005](#)) or the dormancy present in some species ([Potter *et al.* 1984](#), [Mandujano *et al.* 1997](#)). [Silvius \(1995\)](#) observed that *Stenocereus griseus* seeds obtained from bird feces took

between two and five months to germinate, so at least a transient seed bank was possible for the species.

Most of the cactus seeds studied show small size, positive photoblasticism or skotodormancy (seed dormancy after darkness incubation) that allow them to persist in seed banks ([Rojas-Aréchiga & Vázquez-Yanes 2000](#), [Rojas-Aréchiga & Batis 2001](#), [Flores et al. 2006](#), [2011](#), [Ruiz-González et al. 2011](#), [Rojas-Aréchiga et al. 2013](#), [Sosa-Pivatto et al. 2014](#), [Bauk et al. 2017](#), [García-Beltrán et al. 2017](#), [Rojas-Aréchiga & Mandujano-Sánchez 2017](#)). Corresponding to the expected behavior for such seed traits, [Seal et al. \(2009\)](#) published a database with 83 species with seeds that remained viable for more than a year after collection and 65 species with desiccation tolerance. We found one hundred and forty-two cacti species contain seeds capable of remaining viable for at least three months after collection ([Table 1](#)). Even some works where the cactus seeds were stored under controlled conditions have demonstrated that these can germinate after various years ([Flores et al. 2005](#), [Sánchez-Salas et al. 2006](#)) up to a maximum of 10 years ([Alcorn & Martin 1974](#), [Fearn 1977](#), [Trujillo et al. 2014](#)). Few studies have directed their efforts toward evaluating the viability of cactus seeds under natural conditions and their permanence in seed banks ([Álvarez-Espino et al. 2014](#), [Aragón-Gastélum et al. 2018](#)).

Some recent work suggests the possibility of seed banks because of germination trials for various months or years ([Flores et al. 2005](#), [Faife-Cabrera & Toledo-Reina 2007](#), [De la Rosa-Manzano & Briones 2010](#), [Salazar et al. 2013](#), [Jiménez-Sierra & Matías-Palafox 2015](#)) or through the collection and measurement of seed viability ([Cano-Salgado et al. 2012](#)), although in this case it is impossible to determine the age of the collected seeds. Only in 19 species has the existence of a seed bank in the soil been clearly established ([Table 1](#)).

There are few studies that have submitted the seeds to environmental conditions and have examined their viability after different burial periods ([Bowers 2000](#), [2005](#), [Matías-Palafox 2007](#), [Olvera-Carrillo et al. 2009](#), [Cheib & Souza 2012](#), [Goodman et al. 2012](#), [Álvarez-Espino et al. 2014](#), [Ordoñez-Salanueva et al. 2017](#), [Lindow-López et al. 2018a](#), [Aragón-Gastélum et al. 2018](#)) or have assessed the effective seed bank, extracting the seeds from the soil in different times of the year ([Montiel & Montaña 2003](#)). Only the studies of [Bowers \(2005\)](#) and [Ordoñez-Salanueva et al. \(2017\)](#) shows two species (*Mammillaria grahamii* and *Polaskia chende*) capable of possessing a long-term, continuous seed bank (up to at least five years) according to the classification of [Bakker et al. \(1996\)](#).

Serotiny represents an alternative to the traditional seed bank and in many cases is known as an aerial form of seed bank, with the advantage of protecting the seeds in structures that are inaccessible to predators ([Rodríguez-](#)

[Ortega et al. 2006](#)). Serotiny is known as a delayed dispersion mechanism because of the retention of mature seeds in structures of the mother plant for more than a year ([Peters et al. 2009](#)). In the Cactaceae family, serotiny probably is a common phenomenon in species that inhabit several North American deserts like the Mojave and the Sonora ([Martínez-Berdeja et al. 2015](#)). So far 24 species of cacti with a retention of mature seeds have been listed ([Table 1](#)), although around 25 species showed some degree of retention in the work of [Bravo-Hollis & Sánchez-Mejorada \(1991\)](#) according to [Peters et al. \(2009\)](#) but the role that serotiny plays in the population dynamic of the species where it occurs has not been extensively studied.

Seed retention in cacti has been observed in the axils of tubers ([Rodríguez-Ortega et al. 2006](#), [Peters et al. 2009](#)), in chained fruits ([Martínez-Berdeja et al. 2015](#)), in the apex of the stems and in the cephalia ([Bravo-Hollis & Sánchez-Mejorada 1991](#)). The duration of the seeds in the stems has mainly been studied in the *Mammillaria* genus, where it has been demonstrated that the seeds can be retained from one ([Santini & Martorell 2013](#)) to eight years, forming an aerial seed bank ([Boke 1960](#), [Peters et al. 2009](#), [Rodríguez-Ortega et al. 2006](#)).

Temperature and seed germination. The current knowledge about the germination response of cactus seeds concerning different temperatures includes approximately 205 taxa ([Table 1](#)); the majority possesses an optimal germination between 20 - 30 °C ([Zimmer 1968](#), [1982](#), [1998](#), [Rojas-Aréchiga & Vázquez-Yanes 2000](#), [Meiado et al. 2016](#), [Seal et al. 2017](#)). In a few species, greater germination was obtained at 15 °C, including those found in *Parodia aureicentra*, *Pereskia grandifolia* subsp. *grandifolia*, *Pereskia bahiensis*, *Pereskia aculeata*, *Echinopsis schickendantzii*, *Acanthocalycium spiniflorum*, *Ferocactus glaucescens*, *Mammillaria polythele*, *Browningia hertlingiana*, *Cephalocereus senilis*, *Rebutia minuscula*, *Rhipsalis pilocarpa*, *Rhipsalis teres* and *Copiapoa cinerea* var. *haseltoniana* ([Zimmer 1968](#), [1982](#), [1998](#), [Ortega-Baes et al. 2011](#), [Lone et al. 2016](#), [Meiado et al. 2016](#), [Seal et al. 2017](#), [Lindow-López et al. 2018b](#)). Most of the species mentioned above inhabit temperate climates with annual average temperatures between 10.1 and 19.4 °C (<https://es.climate-data.org>), except *Pereskia* spp. and *Rhipsalis* spp. that inhabit warm climates with annual averages between 23.2 and 26.4 °C ([Da Silva 2004](#), <https://es.climate-data.org>).

There are species with different growth forms with seeds that can germinate more than 50 % at temperatures as high as 40 °C. The majority have been reported in columnar species such as: *Pilosocereus gounellei* subsp. *gounellei*, *Pilosocereus pachycladus* subsp. *pernambucoensis* ([Meiado et al. 2016](#)), *Stenocereus thurberi*, *Pachycereus pringlei*,

Pachycereus pecten-aboriginum, *Ferocactus peninsulae* (Yang *et al.* 2003) and *Stenocereus griseus* (Arias & Williams 1978), as well as the globular species *Melocactus curvispinus* subsp. *caesioides* (Arias & Lemus 1984) and the epiphyte *Selenicereus setaceus* (Simão *et al.* 2007). The general tendency however, in cacti is that temperatures equal or higher than 35 °C and lower than 15 °C do not promote germination (Zimmer 1968, 1982, 1998, Rojas-Aréchiga & Vázquez-Yanes 2000, Meiado *et al.* 2016, Seal *et al.* 2017).

Because temperatures continually vary in nature, various studies have compared the germination response of the cactus seeds between constant and alternating temperatures. In such studies, the alternation of temperature can produce three types of responses: (1) lower percentage of germination (Rojas-Aréchiga *et al.* 1997, 1998, 2001, Matias-Palafox 2007, Ortega-Baes & Rojas-Aréchiga 2007, Lindow-López *et al.* 2018b), (2) similar percentage of germination (Godínez-Álvarez & Valiente-Banuet 1998, Ruedas *et al.* 2000, De la Barrera & Nobel 2003, Yang *et al.* 2003, Ramírez-Padilla & Valverde 2005, Matias-Palafox 2007, Sánchez-Soto *et al.* 2010, Ortega-Baes *et al.* 2011, Mazzola *et al.* 2013, Jiménez-Sierra & Matias-Palafox 2015, Kin *et al.* 2015, Lindow-López *et al.* 2018b) or (3) higher percentage of germination (Godínez-Álvarez & Valiente-Banuet 1998).

In some studies, the three types of responses have been observed (Ortega-Baes *et al.* 2010, Meiado *et al.* 2016). From these results, it can be inferred that the germination response concerning the constancy or variation of the temperature becomes species-specific (Ortega-Baes *et al.* 2010) so that a general tendency does not exist in the Cactaceae family such as in other families (Baskin & Baskin 2014).

Some studies of the 1990s (Cancino *et al.* 1993, Nolasco *et al.* 1996, Vega-Villasante *et al.* 1996) demonstrated that exposing *Pachycereus pringlei* and *P. pecten-aboriginum* seeds to temperatures from 55 to 70 °C for various hours, days or even weeks prior to sowing; left some seeds viable. Studies performed on 17 species (Table 1), with seeds that have been exposed to temperatures between 60 - 90 °C (Ruedas *et al.* 2000, Olvera-Carrillo *et al.* 2003, Mandujano *et al.* 2005, Ramírez-Padilla & Valverde 2005, Sánchez-Soto *et al.* 2010, Pérez-Sánchez *et al.* 2011), found four types of response: a) For *Isolatocereus dumortieri* at 70 °C the final germination percentage decreased (Pérez-Sánchez *et al.* 2011); b) For *Mammillaria magnimama*, decreasing only occurred in lots of old seeds (Ruedas *et al.* 2000); c) For several species, the most common response was to show similar germination than in the controls (Ruedas *et al.* 2000, Olvera-Carrillo *et al.* 2003, Ramírez-Padilla & Valverde 2005, Sánchez-Soto *et al.* 2010); d) For several species the exposure to high temperatures exceeded the final

germination of the control (Olvera-Carrillo *et al.* 2003, Sánchez-Soto *et al.* 2010, Pérez-Sánchez *et al.* 2011), demonstrating cactus seed tolerance to high temperatures, at least for short periods. The seed tolerance to extreme temperatures may have evolved as a mechanism to enable persistence in the soil in predominantly desert species with high temperatures on the soil surface (Daws *et al.* 2007).

In recent years, various studies of cactus have directed their efforts to evaluating the possible impact of global warming on the germination of the seeds (Ordoñez-Salanueva *et al.* 2015, Flores *et al.* 2017, Gurvich *et al.* 2017, Seal *et al.* 2017, and Aragón-Gastélum *et al.* 2018), through the assessment of the germination response faced with different temperatures. Other studies, although not directly mentioning global warming, show useful results to evaluate the response of cactus seeds faced with temperatures higher than 30 °C (Kin *et al.* 2015, Meiado *et al.* 2016). However, conclusions are ambiguous.

The results shown by Seal *et al.* (2017) showed a possible explanation for the ambiguity of the response of cacti seeds to high temperatures. That study showed the tendency of species that inhabit temperate environments (extratropical or tropical cactus species that inhabit altitudes > 1,000 m asl) possess optimal germination temperatures generally, between 5 and 10 °C of difference above the average of the period when germination occurs. This study revealed the capability of thermal buffering for the germination that these species possess. Although Seal *et al.* (2017) only included a few species that inhabit warm environment (tropical species that grow between 0 - 1,000 m asl.) the aforementioned species were shown to possess near optimal germination temperatures, generally between 1 and 3 °C of difference or slightly lower than the current average in the rainy period (Seal *et al.* 2017). For these species, the models predict an impact through the increase of the temperatures for the RCP 8.5 scenario (of greater warming) and in some cases until the RCP 2.6 scenario (of lower warming) (Seal *et al.* 2017).

Most germination studies performed on cacti, where different temperatures are compared, generally include species that inhabit temperate environments (Seal *et al.* 2009, 2017). Seal *et al.* (2009) gathered information about germination from approximately a hundred cacti with values between 30 and 41 °C, but only 12.6 % were species from warm climates. From this compilation, it can be deduced that both the cactus species that inhabit temperate climates and those of warm climates generally possess a similar response to this temperature range.

The germination study that included the greatest quantity of tropical cacti species was performed by Meiado *et al.* (2016) with 30 species, mostly endemic of the Caatinga in Brazil. These authors found a germination response higher than 50 % to temperatures of 35 °C and an optimal

temperature around 30 °C, which is a temperature near or slightly higher than the average temperatures reported for that region according to [Da Silva \(2004\)](#). Although it could be expected that the cactus species that inhabit temperate environments have a lower optimal germination temperature than species that inhabit warmer environments, the information available shows that both cacti groups generally possess a similar germination response to 30 °C ([Figure 4](#)).

Figure 4. Seed germination responses at 30 °C between cacti from temperate and warm climates, based on 43 data from temperate and 34 from warm climate.

This similar response implies that the cactus species of temperate climates possess the greatest capacity for thermal buffering of germination, as the results of [Seal et al. \(2017\)](#) suggest. The warm climate species, being within the limit or very close to the optimal germination temperature will possess a lower capacity for buffering and in consequence it will possibly be these cacti that are the most vulnerable to the effects of the expected climatic change.

Light and seed germination. The germination response considering conditions of light and darkness has been assessed in 275 cactus species ([Table 1](#)). So far no negative photoblastic seeds have been found in cacti and 80.9 % have presented positively photoblastic seeds ([Matías-Palafox & Jiménez-Sierra 2006](#), [Flores et al. 2006, 2011](#), [Cheib & Souza 2012](#), [Delgado-Sánchez et al. 2013](#), [Rojas-Aréchiga et al. 2013](#), [Meiado et al. 2016](#)). Among the cacti with indifferent or neutral photoblastic seeds, 34 taxa have been reported in the Cactoideae subfamily that represent only 13.8 % of its species, however, 75 % of the 24 taxa studied of the subfamilies Opuntioideae, Pereskioideae and Maihuenioideae, have indifferent photoblastic seeds, which suggests that this character is ancestral in Cactaceae.

Several studies indicate a strong influence from phylogeny in the photoblastic response within the family

([Flores et al. 2011](#), [Rojas-Aréchiga et al. 2013](#), [Meiado et al. 2016](#)). [Rojas-Aréchiga et al. \(2013\)](#) found a phylogenetic influence between various species of the Cactaceae tribe, while [Flores et al. \(2011\)](#) found that the Cactaceae, Pachycereeae and Trichocereae tribes include species with greater germination response to light than in Notocactaceae. On the other hand, [Meiado et al. \(2016\)](#) found that the phylogenetic origin was the only variable influencing the germination response to light in 30 cactus species of Brazil, and only the species of Pereskioideae and Opuntioideae showed an indifferent photoblastic response, while all the Cactoideae species were positively photoblastic.

The seed response to light in Cactaceae has traditionally been evaluated in association with the form of life ([Rojas-Aréchiga et al. 1997](#), [Ortega-Baes et al. 2010](#), [Flores et al. 2011](#)), size and seed mass ([Flores et al. 2006](#), [Ortega-Baes et al. 2010](#), [Rojas-Aréchiga et al. 2013](#)), as well as to different wavelengths or solar irradiation ([Alcorn & Kurtz 1959](#), [McDonough 1964](#), [Zimmer 1969a, 1973](#), [Rojas-Aréchiga et al. 1997](#), [Nolasco et al. 1997](#), [Benítez-Rodríguez et al. 2004](#), [Olvera-Carrillo et al. 2009](#), [Meiado et al. 2010](#), [Zerpa-Catanho et al. 2019](#)) and concentrations of gibberellins ([Alcorn & Kurtz 1959](#), [McDonough 1964](#), [Rojas-Aréchiga et al. 2001](#), [Ortega-Baes & Rojas-Aréchiga 2007](#), [Rojas-Aréchiga et al. 2011](#), [Mascot-Gómez et al. 2020](#)). Some preliminary studies suggested that an association between the life-form and the photoblasticism could exist ([Rojas-Aréchiga et al. 1997](#), [Rojas-Aréchiga & Vázquez-Yanes 2000](#)). Other research has subsequently opposed that hypothesis ([Ortega-Baes et al. 2010](#), [Flores et al. 2011](#), [Meiado et al. 2016](#)). Moreover, [Flores et al. \(2011\)](#) found a negative relationship between the germination response to light and the height of the plant, probably because tall cacti produce larger seedlings than short cacti ([Flores et al. 2011](#)). Similar readings can be obtained from the data of [Rojas-Aréchiga et al. \(1997\)](#).

Several cactus studies have shown a relationship between seed size and the requirement of light to germinate ([Maiti et al. 2003](#), [Sánchez et al. 2015](#)). Both [Flores et al. \(2006\)](#) and [Ortega-Baes et al. \(2010\)](#) considered that positive photoblasticism in the 37 cactus species they studied could be related to the small size of their seeds and their potential for building soil seed banks. Similarly, [Flores et al. \(2011\)](#) obtained a negative relationship between seed mass and the dependence on light to germinate; nevertheless, in species without dormancy this relationship was marginal or non-significant, which the authors attributed to the differences that were presented between dormant and non-dormant seeds or to the size of the sample.

Some previous studies of around 80 species of cacti ([Rojas-Aréchiga et al. 2013](#), [Meiado et al. 2016](#)) found no relationship between the requirement of light for

germinating and seed size. Probably these results with a certain ambiguity could be explained by the small size and mass that cacti show in general and because the role of the phylogenetic relationships (as already referred to) can distort the interpretation of the data.

The influence of gibberellic acid (AG_3) as a light substitute has been studied in cacti since [Alcorn & Kurtz \(1959\)](#) found that a fraction of positively photoblastic seeds of *Carnegiea gigantea* were stimulated in darkness with the addition of AG_3 . Subsequently, [McDonough \(1964\)](#) confirmed the aforementioned results and also included *Stenocereus thurberi* with similar results. Similar findings have been found in several cactus genera by [Brencher *et al.* \(1978\)](#), [Arias & Williams \(1978\)](#), [Zimmer & Büttner \(1982\)](#) and [López-Gómez & Sánchez-Romero \(1989\)](#). Nevertheless, even though the role of AG_3 has not been subsequently studied ([Ortega-Baes & Rojas-Aréchiga 2007](#), [Rojas-Aréchiga *et al.* 2011](#)), the works that refer to it in cacti have not found a stimulation of germination under darkness ([Williams & Arias 1978](#), [Arias & Lemus 1984](#), [Rojas-Aréchiga *et al.* 2001, 2011](#), [Olvera-Carrillo *et al.* 2003](#), [Ortega-Baes & Rojas-Aréchiga 2007](#), [Rojas-Aréchiga 2008](#), [Mascot-Gómez *et al.* 2020](#)). Even when the AG_3 effect as a promoter of germination under white light has been evaluated, in the majority of the studies an enhancer effect on different concentrations in intact seeds has not been found ([Mandujano *et al.* 2007](#), [Ortega-Baes & Rojas-Aréchiga 2007](#), [Rojas-Aréchiga 2008](#), [Olvera-Carrillo *et al.* 2009](#), [Amador-Alfárez *et al.* 2013](#), [Loustalot *et al.* 2014](#), [Rodríguez-Ruiz *et al.* 2018](#), [Gonzalez-Cortés *et al.* 2018](#)). The difficulty in clearly establishing the role of AG_3 as a germination promoter for Cactaceae could be related to the fact that in the studies homogeneity has not been maintained in the following parameters: light conditions, age of the seeds, types of dormancy, time and mode of application of the AG_3 concentrations, among others. These differences among parameters can include changes in the behavior of seeds' germination in the species and thus hinder the comparisons among the performed studies ([Rojas-Aréchiga 2008](#)). Nevertheless, possibly AG_3 does not possess such an important role as a germination promoter in cacti seeds, such as [Rojas-Aréchiga \(2008\)](#) and [Rojas-Aréchiga *et al.* \(2011\)](#) refer to.

Water and saline stress and germination. In general, seeds of most plants studied stop germinating at water potentials between -0.5 and -2.0 MPa ([Dürr *et al.* 2015](#), [Tribouillois *et al.* 2016](#)). These potentials are experimentally achieved with the use of different concentrations of three main substances: mannitol ([Maldonado *et al.* 2002](#), [Guerrero *et al.* 2016](#)), polyethylene glycol (PEG) ([Yang *et al.* 2010](#), [Luna & Chamorro 2016](#), [Pantané *et al.* 2016](#)) and NaCl ([Zhang *et al.* 2010](#), [Gorai *et al.* 2014](#), [El-Keblawy *et al.* 2016](#)).

In cacti, only four studies have assessed the germination at water potentials lower than -1.0 MPa ([Martins *et al.* 2012](#), [Guerrero *et al.* 2016](#), [Aragón & Lasso 2018](#), [Zerpa-Catanho *et al.* 2019](#)), maybe because in most cases the species do not achieve germination at such potentials ([Flores & Briones 2001](#), [Ramírez-Padilla & Valverde 2005](#), [De La Rosa-Manzano & Briones 2010](#), [Guillén *et al.* 2011, 2015](#), [Rodríguez-Morales *et al.* 2013](#), [Flores *et al.* 2017](#), [Bispo *et al.* 2018](#)). The studies in cacti reveal that germination mainly occurs at water potentials between 0 and -0.6 MPa ([Figure 5](#)) and at temperatures of 25 to 30 °C ([Vega-Villasante *et al.* 1996](#), [Kin *et al.* 2015](#), [Gurvich *et al.* 2017](#), [Flores *et al.* 2017](#), [Bispo *et al.* 2018](#)).

Figure 5. Seed germination response under water stress. Data are based on 28 cactus species studied.

In some species, it has been observed that the water potentials of -0.2 MPa ([Guillén *et al.* 2009, 2015](#), [Gurvich *et al.* 2017](#), [Flores *et al.* 2017](#)) or even -0.4 MPa ([Flores & Briones 2001](#), [Guillén *et al.* 2011, 2015](#), [Flores *et al.* 2017](#)) or -0.6 MPa ([Flores & Briones 2001](#)) favor germination. Nevertheless, generally water potentials lower than -0.6 MPa notably decrease the percentage of germination ([Meiado *et al.* 2010](#), [Guillén *et al.* 2011, 2015](#), [Gurvich *et al.* 2017](#), [Flores *et al.* 2017](#), [Bispo *et al.* 2018](#)). Thus, cactus seeds adapted to germinate at high soil moisture, but not necessarily at the soil field water capacity, would have an advantage germinating in arid environments. However, this may have a trade-off between germination at relatively low soil water potential and seedling establishment because seedlings generally require higher soil water potential to compensate for transpiration ([Flores & Briones 2001](#)).

In addition, in the species where the effect of water potential and temperature has been combined, normally at 25 °C, better results have been obtained than at temperatures of 30 °C or higher ([Oliveira *et al.* 2017](#), [Gurvich *et al.* 2017](#), [Flores *et al.* 2017](#), [Bauk *et al.* 2017](#)). For primarily arid and semiarid species with small seeds, as

occurs in Cactaceae, the preference for germinating in humid soil can be an advantage that permits to reduce seedling mortality when faced with transient rain followed by prolonged droughts. These conditions could favor germination but could also result in death of seedlings through subsequent drought (Venable & Lawlor 1980).

Approximately a dozen studies (Table 1) have evaluated the germination response of cacti faced with conditions of salinity (Romero-Schmidt *et al.* 1992, Vega-Villasante *et al.* 1996, Nolasco *et al.* 1996, Meiado *et al.* 2010, Martins *et al.* 2012, Ortiz *et al.* 2014, Lima & Meiado 2017, Podda *et al.* 2017, Bispo *et al.* 2018). A germination response in the range of 0 to -0.6 MPa with NaCl is similar to the results obtained with PEG, even though there are species where the germination percentages with NaCl are higher than in a similar osmotic potential of PEG (Meiado *et al.* 2010, Bispo *et al.* 2018). In other cases, the seeds germinate better in PEG (Martins *et al.* 2012). On the other hand, there is very little information about the application of the hydrotime model to evaluate the effects of water or saline stress on germination speed (or germination time) in cactus species (see: Simão *et al.* 2010). Germination time is an important feature of plant life history that determines germination capacity under dry conditions and recruitment time (Sánchez-Soto *et al.* 2005, Donohue *et al.* 2010, Dürr *et al.* 2015).

Hydration-dehydration cycles and germination. Two pioneering studies of Dubrovsky (1996, 1998) demonstrated the utility of the hydration-dehydration cycles in increasing the speed and final percentage of germination in four cactus species. However, still there are a few cactus studies (Table 1) that address these treatments as germination promoters (Sánchez-Soto *et al.* 2005, Rito *et al.* 2009, Santini & Martorell 2013, López-Urrutia *et al.* 2014, Contreras-Quiroz *et al.* 2016a, 2016b, Santini *et al.* 2017, Lima & Meiado 2017, 2018, Sánchez *et al.* 2018), although hydration-dehydration cycles have been and are widely used in innumerable species worldwide (Sánchez & Pernús 2018). The evaluation of this type of pre-germinative treatment in cactus seeds should be tested more in the future, because the seeds are subject to hydration-dehydration cycles in nature, which can influence not only germination but also aspects related to conditional or cyclical dormancy, seed longevity and seedling survival, among others (Baskin & Baskin 2014). Similarly, water treatments (or cycles) could be implemented as ecotechnologies in ecological restoration projects and in conservation programs of germplasm (Sánchez & Pernús 2018).

The ability of seeds to maintain physiological changes produced during hydration, such as differential protein expression, through discontinuous dehydration periods (López-Urrutia *et al.* 2014), has been called “hydration

memory” (Dubrovsky 1996). Contreras-Quiroz *et al.* (2016b) evaluated this phenomenon in cacti from a semi-desert area in northeastern Mexico, and from a sub-humid area of central Argentina, and suggested that the presence of hydration memory in the seeds of Cactaceae depends on the climate and the microenvironment where the cacti occur, indicating that environmental conditions imposed on the parental plants influence the germination responses of the cacti seeds when subjected to discontinuous hydration. This hypothesis was corroborated by Lima & Meiado (2017) who found that the seeds of the same species (*Pilosocereus catingicola* subsp. *salvadorensis*) collected from populations located in different ecosystems have different germination responses after passage through discontinuous hydration and HD cycles. This finding gives evidence that “hydration memory” provides greater tolerance to environmental stresses but with different responses among populations.

Cactus seed dispersal. Rojas-Aréchiga & Vázquez-Yanes (2000) reported six studies about cactus seed dispersion through endozoochory. In the last two decades, a few more than a dozen species have been studied regarding this type of interaction and its effect on germination (Godínez-Álvarez & Valiente-Banuet 2000, Montiel & Montaña 2000, Godínez-Álvarez *et al.* 2002, Naranjo *et al.* 2003, Baraza & Valiente-Banuet 2008, Pérez-Villafaña & Valiente-Banuet 2009, Casado & Soriano 2010, Fonseca *et al.* 2012, Gomes *et al.* 2014, Nascimento *et al.* 2015, Lasso & Barrientos 2015, Vázquez-Castillo *et al.* 2018, Santos *et al.* 2019). Other types of cactus seed dispersal referred to by Rojas-Aréchiga & Vázquez-Yanes (2000) as synzoochory (Munguía-Rosas *et al.* 2009, Fonseca *et al.* 2012), epizoochory (Lasso & Barrientos 2015) and hydrochory (Lenzi *et al.* 2012, Sánchez-Salas *et al.* 2012, Sánchez-Salas *et al.* 2013, Romero-Méndez *et al.* 2018) have been reported in a small number of studies.

In vitro propagation. Although in a great number of cactus species high percentages of germination have been found (Zimmer 1969b, 1998, Meiado *et al.* 2016), various studies have explored *in vitro* germination as an alternative of propagation (Salas-Cruz *et al.* 2011, Xavier & Jasmim 2015, López-Escamilla *et al.* 2016, Cortés-Olmos *et al.* 2018, Ramírez-González *et al.* 2019). Studies of this type have been performed on more than 40 cactus species (Table 1), with good results in germination and later seedling growth. The *in vitro* germination with enriched resources could be a necessary and essential method in the propagation of rare species with few seeds or with a low percentage of germination, both for preservation of the species or another purpose.

Seed storage at low temperatures. Although in the present review the seed desiccation tolerance and the longevity of cactus seeds have been commented on, few studies have evaluated the seed preservation in controlled conditions. Only in 20 species has a preservation at -0°C been evaluated, up to a maximum of 13 months (Nolasco *et al.* 1996, Vega-Villasante *et al.* 1996, Veiga-Barbosa *et al.* 2010, Goodman *et al.* 2012, Marchi *et al.* 2013, Salazar *et al.* 2013, Bárbara *et al.* 2015, Civatti *et al.* 2015, Dos Santos *et al.* 2018). Studies of this type are necessary when faced with the increasing threats that confront the family (Goettsch *et al.* 2015) and the need to search for secure alternatives for long-term preservation.

Concluding remarks. In the last two decades, the number of studies related to cactus germination has increased exponentially, with Mexico, Brazil and Argentina being the three countries leading this scientific field. Seed germination has been evaluated in less than 50 % of the genera. Only physiological dormancy has been found in the family. Terrestrial and aerial cactus seed banks have been little studied but are perhaps common and play an important role in the population dynamics of the species. The two most studied research topics on cactus seed germination are photoblasticism and the response to temperature. No cactus with a negatively photoblastic response has been found, and most of the species of the subfamily *Cactoideae* have positively photoblastic seeds. The optimal temperature range for cacti is found between 20 and 30 $^{\circ}\text{C}$. However, there are still genera with more than 20 species that have not been studied such as *Matucana*, and others genera like *Frailea*, *Rebutia*, and *Selenicereus* where only one species has been studied.

There is not enough evidence on seed traits and cactus seed germination, but seed shape has been related with seed dispersal in that the hat-shaped seeds have been associated with hydrochory. The relationships between cactus seed germination and phylogeny depends on the species studied and the tribe. The germination types or variants do not appear to be related with seed germination percentage. Cactus seeds are adapted to germinate at high soil moisture, but not necessarily at the soil field water capacity. Seeds germinate at saline potentials between 0 and -0.6 MPa. Hydration-dehydration cycles or hydration memory increase the speed and the final percentage of germination in some species; the presence of hydration memory in the seeds of *Cactaceae* depends on the climate and the microenvironment where the cacti occur.

Preserving cactus seeds in banks is an encouraging concept because their longevity in general surpasses six months, and a high percentage of species possess seeds that maintain their viability for periods between one to two years without extreme storage requirements. Similarly, studies

about desiccation tolerance and cryopreservation that have been performed demonstrate their feasibility. In the future, it would be useful if more cactus germination studies included measurement of seed traits, such as length, width and compression, total dry mass, mass of the covering of the embryo, moisture content, as well as imbibition rate.

Acknowledgements

The present study relied on support from the National Botanical Garden, the Cuban Botanical Society and Planta! The first author is grateful for the support received from R. Rankin, L. R. González-Torres, A. C. González, J.L. Aragón-Gastelum, M. V. Meiado and D. Franco for the literature they sent. We appreciate the help of A. Ruiz Fleitas with the seed drawings. We also thank the journal editors and two anonymous reviewers whose work contributed to improving the paper.

Literature cited

- Alcorn SM, Kurtz Jr EB. 1959. Some factors affecting the germination of seed of the saguaro cactus (*Carnegiea gigantea*). *American Journal of Botany* **46**: 526-529. DOI: <https://doi.org/10.2307/2439624>
- Alcorn SM, Martin SC. 1974. *Cereus giganteus* Engelm. In: Schopmeyer C, ed. *Seeds of Woody Plants in the United States*. Washington, DC: Forest Service US. Department of Agriculture, 313-314.
- Almeida OJG, Souza LAd, Moscheta IS. 2009. Morfo-anatomía da plântula de indivíduos somaclones de *Cereus hildmannianus* Schumann (Cactaceae). *Boletín de la Sociedad Latinoamericana y del Caribe de Cactáceas y otras Suculentas* **6**: 29-35.
- Álvarez-Espino R, Godínez-Álvarez H, Torre-Almaráz R. 2014. Seed banking in the columnar cactus *Stenocereus stellatus*: distribution, density and longevity of seeds. *Seed Science Research* **24**: 315-320. DOI: <https://doi.org/10.1017/S0960258514000324>
- Amador-Alfárez KA, Díaz-González J, Loza-Cornejo S, Bivián-Castro EY. 2013. Efecto de diferentes reguladores de crecimiento vegetal sobre la germinación de semillas y desarrollo de plántulas de dos especies de *Ferocactus* (Cactaceae). *Polibotánica* **35**: 109-131.
- Anderson EF. 2001. *The Cactus Family*. Portland, Oregon: Timber Press, Inc. ISBN 10: 0-88192-498-9; ISBN-13: 978-0881924985
- Aragón L, Lasso E. 2018. How do young cacti (seeds and seedlings) from tropical xeric environments cope with extended drought periods? *Journal of Arid Environments* **154**: 1-7. DOI: <https://doi.org/10.1016/j.jaridenv.2018.03.009>

- Aragón-Gastélum JL, Reyes-Olivas Á, Sánchez-Soto BH, Casillas-Álvarez P, Flores J. 2013. Vivipary in *Ferocactus herrerae* (Cactaceae) in Northern Sinaloa, México. *Bradleya* **31**: 44-52. DOI: <https://doi.org/10.25223/brad.n31.2013.a7>
- Aragón-Gastélum JL, Flores J, Yáñez-Espinosa L, Reyes-Olivas A, Rodas-Ortiz JP, Robles-Díaz E, González FJ. 2017. Advantages of vivipary in *Echinocactus platyacanthus*, an endemic and protected Mexican cactus species. *Journal of Arid Environments* **141**: 56-59. DOI: <http://dx.doi.org/10.1016/j.jaridenv.2017.01.012>
- Aragón-Gastélum JL, Flores J, Jurado E, Ramírez-Tobías HM, Robles-Díaz E, Rodas-Ortiz JP, Yáñez-Espinosa L. 2018. Potential impact of global warming on seed bank, dormancy and germination of three succulent species from the Chihuahuan Desert. *Seed Science Research* **28**: 312-318. DOI: <https://doi.org/10.1017/S0960258518000302>
- Areces-Mallea A. 2003. *Leptocereus* (A. Berger) Britton and Rose: a monographic study of a West Indian genus of Cactaceae (Cactoideae). PhD Thesis, City University of New York.
- Arias I, Williams PM. 1978. Physio-ecological studies of plant species from the arid and semi-arid regions of Venezuela. *Acta Científica Venezolana* **29**: 384-388.
- Arias I, Lemus L. 1984. Interaction of light, temperature and plant hormones in the germination of seeds of *Melocactus caesius* Went. (Cactaceae). *Acta Científica Venezolana* **35**: 151-155.
- Arias S, Terrazas T. 2004. Seed morphology and variation in the genus *Pachycereus* (Cactaceae). *Journal of Plant Research* **117**: 277-289. DOI: <https://doi.org/10.1007/s10265-004-0156-4>
- Arias S, Gama-López S, Guzmán-Cruz LU, Vázquez-Benítez B. 2012. *Flora del Valle de Tehuacán-Cuicatlán. Cactaceae*. Ciudad de México: Universidad Nacional Autónoma de México. ISBN: 978-607-02-3079-0
- Arroyo-Cosultchi G, Terrazas T, Arias S, Arreola-Nava HJ. 2006. The systematic significance of seed morphology in *Stenocereus* (Cactaceae). *Taxon* **55**: 983-992. DOI: <https://doi.org/10.2307/25065693>
- Arroyo-Cosultchi G, Terrazas T, Arias S, López-Mata L. 2007. Morfología de las semillas en *Neobuxbaumia* (Cactaceae). *Boletín de la Sociedad Botánica de México* **81**: 17-25. DOI: <https://doi.org/10.17129/botsoci.1762>
- Bakker JP, Poschod P, Strykstra RJ, Bekker RM, Thompson K. 1996. Seed banks and seed dispersal: Important topics in restoration ecology. *Acta Botanica Neerlandica* **45**: 461-490. DOI: <https://doi.org/10.1111/j.1438-8677.1996.tb00806.x>
- Baraza E, Valiente-Banuet A. 2008. Seed dispersal by domestic goats in a semiarid thornscrub of Mexico. *Journal of Arid Environments* **72**: 1973-1976. DOI: <https://doi.org/10.1016/j.jaridenv.2008.04.004>
- Bárbara EPS, Silva AA, Souza MMOR, Gurgel ZER, Marchi MNG, Bellintani MC. 2015. Germinação e criopreservação de sementes de cactos nativos da Bahia. *Gaia Scientia* **9**: 91-96.
- Barrios D, González-Torres LR, García-Beltrán JA. 2012. Vivipary in Cuban cacti: a pioneer study in *Leptocereus scopulophilus*. *Bradleya* **30**: 147-150. DOI: <https://doi.org/10.25223/brad.n30.2012.a17>
- Barrios D, Flores J, González-Torres LR, Palmarola A. 2015. The role of mucilage in the germination of *Leptocereus scopulophilus* (Cactaceae) seeds from Pan de Matanzas, Cuba. *Botany* **93**: 251-255. DOI: <https://doi.org/10.1139/cjb-2014-0242>
- Barthlott W, Hunt D. 1993. Cactaceae. In: Kubitzki K, Rohrer JG, Bittrich V. eds. *The Families and Genera of Vascular Plants*. Berlin Heidelberg: Springer-Verlag, 161-197. ISBN: 978-3-642-08141-5, 978-3-662-02899-5; DOI: https://doi.org/10.1007/978-3-662-02899-5_17
- Barthlott W, Hunt D. 2000. *Seed diversity in the Cactaceae subfamily Cactoideae*. Milborne, Port: DH Books. ISBN: 97809538134 07
- Barthlott W, Porembski S. 1996. Ecology and morphology of *Blossfeldia liliputana* (Cactaceae): a poikilohydric and almost astomate succulent. *Botanica Acta* **109**: 161-166. DOI: <https://doi.org/10.1111/j.1438-8677.1996.tb00556.x>
- Barthlott W, Porembski S, Kluge M, Hopke J, Schmidt L. 1997. *Selenicereus wittii* (Cactaceae): an epiphyte adapted to Amazonian Igapó inundation forests. *Plant Systematics and Evolution* **206**: 175-185. DOI: <https://doi.org/10.1007/BF00987947>
- Baskin CC, Baskin JM. 2014. *Seeds: Ecology, biogeography, and evolution of dormancy and germination*. San Diego: Elsevier Sciences. ISBN 13: 978-0-12-416677-6
- Bauk K, Flores J, Ferrero C, Pérez-Sánchez R, Las Peñas ML, Gurruchaga DE. 2017. Germination characteristics of *Gymnocalycium monvillei* (Cactaceae) along its entire altitudinal range. *Botany* **95**: 419-428. DOI: <https://doi.org/10.1139/cjb-2016-0154>
- Benítez-Rodríguez LJ, Orozco-Segovia A, Rojas-Aréchiga M. 2004. Light effect on seed germination of four *Mammillaria* species from the Tehuacán-Cuicatlán valley, Central Mexico. *The Southwestern Naturalist* **49**: 11-17.
- Bispo JP, Meiado MV, Siqueira-Filho JA. 2018. Seed germination of three endangered subspecies of *Discocactus Pfeiff* (Cactaceae) in response to environmental factors. *Journal of Seed Science* **40**: 253-262. DOI: <http://dx.doi.org/10.1590/2317-1545v40n3183036>

- Boke NH. 1960. Anatomy and development in *Solisia*. *American Journal of Botany* **47**: 59-65. DOI: <https://doi.org/10.2307/2439495>
- Bowers JE. 2000. Does *Ferocactus wislizenii* (Cactaceae) have a between-year seed bank? *Journal of Arid Environments* **45**: 197-205. DOI: <https://doi.org/10.1006/jare.2000.0642>
- Bowers JE. 2005. New evidence for persistent or transient seed banks in three Sonoran Desert cacti. *The Southwestern Naturalist* **50**: 482-487. DOI: [https://doi.org/10.1894/0038-4909\(2005\)050\[0482:NEFPOT\]2.0.CO;2](https://doi.org/10.1894/0038-4909(2005)050[0482:NEFPOT]2.0.CO;2)
- Bravo-Hollis H. 1978. *Las Cactáceas de México*. México DF: Universidad Nacional Autónoma de México.
- Bravo-Hollis H, Sánchez-Mejorada H. 1991. *Las Cactáceas de México*. México DF: Universidad Nacional Autónoma de México.
- Bregman R, Bouman F. 1983. Seed germination in Cactaceae. *Botanical Journal of Linnean Society* **86**: 357-374. DOI: <https://doi.org/10.1111/j.1095-8339.1983.tb00977.x>
- Bregman R, Graven P. 1997. Subcuticular secretion by cactus seeds improves germination by means of rapid uptake and distribution of water. *Annals of Botany* **80**: 525-531. DOI: <https://doi.org/10.1006/anbo.1997.0483>
- Brencher W, Stange L, Zimmer K. 1978. Ersatz des Lichts bei der Keimung von Kakteensamen durch Gibberellinsäure. *Die Gartenbauwissenschaft* **43**: 91-94.
- Cancino J, León de la Luz J, Coria R, Romero H. 1993. Effect of heat treatment on germination of seeds of cardón [*Pachycereus pringlei* (S. Wats.) Britt. & Rose, Cactaceae]. *Journal of the Arizona-Nevada Academy of Science* **27**: 49-54.
- Cano-Salgado A, Zavala-Hurtado JA, Orozco-Segovia A, Valverde-Valdés MT, Pérez-Rodríguez P. 2012. Composición y abundancia del banco de semillas en una región semiárida del trópico mexicano: patrones de variación espacial y temporal. *Revista Mexicana de Biodiversidad* **83**: 437-446. DOI: <http://dx.doi.org/10.22201/ib.20078706e.2012.2.938>
- Carbajal MCN, Saldivar S, Eliosa HR. 2010. Efecto de la escarificación y de la edad de semillas en la germinación de *Mammillaria mystax*. *Zonas Áridas* **14**: 196-205. DOI: <https://doi.org/10.21704/za.v14i1.161>
- Casado R, Soriano PJ. 2010. Fructificación, frugivoría y dispersión en el cactus globular *Melocactus schatzlii* en el enclave semiárido de Lagunillas, Mérida, Venezuela. *Ecotrópicos* **23**: 18-36.
- Cheib AL, Souza Q. 2012. Longevity and germination ecology of seeds of endemic Cactaceae species from high-altitude sites in south-eastern Brazil. *Seed Science Research* **22**: 45-53. DOI: <https://doi.org/10.1017/S0960258511000298>
- Civatti LM, Marchi MNG, Bellintani MC. 2015. Cryopreservation of cacti seeds of three ornamental species endemic to the state of Bahia, Brazil. *Seed Science and Technology* **43**: 284-290. DOI: <http://doi.org/10.15258/sst.2015.43.2.08>
- Contreras-Quiroz MDR, Pando Moreno M, Flores J, Jurado E. 2016a. Effects of wetting and drying cycles on the germination of nine species of the Chihuahuan Desert. *Botanical Sciences* **94**: 221-228. DOI: <http://dx.doi.org/10.17129/botsci.457>
- Contreras-Quiroz M, Pando-Moreno M, Jurado E, Flores J, Bauk K, Gurvich DE. 2016b. Is seed hydration memory dependent on climate? Testing this hypothesis with Mexican and Argentinian cacti species. *Journal of Arid Environments* **130**: 94-97. DOI: <http://dx.doi.org/10.1016/j.jaridenv.2016.03.001>
- Cortés-Olmos C, Gurrea-Ysasi G, Prohens J, Rodríguez-Burruezo A, Fita A. 2018. *In vitro* germination and growth protocols of the ornamental *Lophophora williamsii* (Lem.) Coult. as a tool for protecting endangered wild populations. *Scientia Horticulturae* **237**: 120-127. DOI: <https://doi.org/10.1016/j.scienta.2018.03.064>
- Côrtes JE, Vasconcellos-Neto J, Garcia MA, Teixeira de Souza AL. 1994. Saurocory in *Melocactus violaceus* (Cactaceae). *Biotropica* **26**: 295-301. DOI: <https://doi.org/10.2307/2388850>
- Cota-Sánchez JH. 2004. Vivipary in the Cactaceae: Its taxonomic occurrence and biological significance. *Flora* **199**: 481-490. DOI: <https://doi.org/10.1078/0367-2530-00175>
- Cota-Sánchez JH, Reyes-Oliva A, Sánchez-Soto B. 2007. Vivipary in coastal cacti: a potential reproductive strategy in halophytic environments. *American Journal of Botany* **94**: 1577-1581. DOI: <https://doi.org/10.3732/ajb.94.9.1577>
- Cota-Sánchez JH, Reyes-Olivas Á, Abreu DD. 2011. Vivipary in the cactus family: A reply to Ortega-Baes' et al. evaluation of 25 species from northwestern Argentina. *Journal of Arid Environments* **75**: 878-880. DOI: <https://doi.org/10.1016/j.jaridenv.2011.04.015>
- Da Silva VPR. 2004. On climate variability in Northeast of Brazil. *Journal of Arid Environments* **58**: 575-596. DOI: <http://dx.doi.org/10.1016/j.jaridenv.2003.12.002>
- Daws MI, Kabadajic A, Manger K, Kranner I. 2007. Extreme thermo-tolerance in seeds of desert succulents is related to maximum annual temperature. *South African Journal of Botany* **73**: 262-265. DOI: <https://doi.org/10.1016/j.sajb.2006.12.004>
- Dehan B, Pérez H. 2005. Preliminary study shows germination of Caribbean applecactus (*Harrisia fragrans*) improved with acid scarification and giberellic acid. *Native Plants Journal* **6**: 91-96. DOI: <https://doi.org/10.1353/npj.2005.0017>

- De la Barrera E, Nobel PS. 2003. Physiological ecology of seed germination for the columnar cactus *Stenocereus queretaroensis*. *Journal of Arid Environments* **53**: 297-306. DOI: <https://doi.org/10.1006/jare.2002.1050>
- De La Rosa-Manzano E, Briones O. 2010. Germination response of the epiphytic cactus *Rhipsalis baccifera* (J. S. Miller) stearn to different light conditions and water availability. *International Journal of Plant Science* **171**: 267-274. DOI: <https://doi.org/10.1086/650159>
- de Viana ML. 1999. Seed production and seed bank of *Trichocereus pasacana* (Cactaceae) in northwestern Argentina. *Tropical Ecology* **40**: 79-84.
- Del Castillo RF. 1988. Fenología y remoción de semillas en *Ferocactus histrix*. *Cactáceas y Suculentas Mexicanas* **33**: 5-14.
- Delgado-Sánchez P, Ortega-Amaro MA, Rodríguez-Hernández AA, Jiménez-Bremont JF, Flores J. 2010. Further evidence from the effect of fungi on breaking *Opuntia* seed dormancy. *Plant Signaling & Behavior* **5**: 1229-1230. DOI: <https://doi.org/10.4161/psb.5.10.12835>
- Delgado-Sánchez P, Ortega-Amaro MA, Jiménez-Bremont JF, Flores J. 2011. Are fungi important for breaking seed dormancy in desert species? Experimental evidence in *Opuntia streptacantha* (Cactaceae). *Plant Biology* **13**: 154-159. DOI: <https://doi.org/10.1111/j.1438-8677.2010.00333.x>
- Delgado-Sánchez P, Jiménez-Bremont JF, Guerrero-González ML, Flores J. 2013. Effect of fungi and light on seed germination of three *Opuntia* species from semiarid lands of central Mexico. *Journal of Plant Research* **126**: 643-664. DOI: <https://doi.org/10.1007/s10265-013-0558-2>
- Donohue K, de Casas RR, Burghardt L, Kovach K, Willis CG. 2010. Germination, postgermination adaptation, and species ecological ranges. *Annual Review of Ecology and Systematics* **41**: 293-319. DOI: <https://doi.org/10.1146/annurev-ecolsys-102209-144715>
- Dos Santos AP, Hassemer G, Meiado MV. 2018. Seed storage of Brazilian cacti species in different threat categories. *Plant Species Biology* **33**: 203-211. DOI: <https://doi.org/10.1111/1442-1984.12208>
- Dubrovsky JG. 1996. Seed hydration memory in Sonoran Desert cacti and its ecological implication. *American Journal of Botany* **83**: 624-632. DOI: <https://doi.org/10.2307/2445922>
- Dubrovsky JG. 1998. Discontinuous hydration as a facultative requirement for seed germination in two cactus species of the Sonoran Desert. *Journal of the Torrey Botanical Society* **125**: 33-39. DOI: <https://doi.org/10.2307/2997229>
- Dürr C, Dickie JB, Yang X-Y, Pritchard HW. 2015. Ranges of critical temperature and water potential values for the germination of species worldwide: Contribution to a seed trait database. *Agricultural and Forest Meteorology* **200**: 222-232. DOI: <http://dx.doi.org/10.1016/j.agrformet.2014.09.024>
- El-Keblawy A, Gairola S, Bhatt A. 2016. Maternal salinity environment affects salt tolerance during germination in *Anabasis setifera*: A facultative desert halophyte. *Journal of Arid Land* **8**: 254-263. DOI: <https://doi.org/10.1007/s40333-015-0023-2>
- Faife-Cabrera M, Toledo-Reina SP. 2007. Viabilidad de semillas de *Melocactus gutartii* León conservadas por seis y doce meses. *Centro Agrícola* **34**: 47-50.
- Fearn B. 1977. An investigation into the effect of age on the germination potential of seeds of 600 species of cacti, together with a note on the viability of *Lithops* seeds. *Excelsa* **7**: 103-108.
- Finch-Savage WE, Leubner-Metzger G. 2006. Seed dormancy and the control of germination. *New Phytologist* **171**: 501-523. DOI: <https://doi.org/10.1111/j.1469-8137.2006.01787.x>
- Fleming TH, Maurice S, Buchmann SL, Tuttle MD. 1994. Reproductive biology and relative male and female fitness in a trioecious cactus, *Pachycereus pringlei* (Cactaceae). *American Journal of Botany* **81**: 858-867. DOI: <http://dx.doi.org/10.1002/j.1537-2197.1994.tb15567.x>
- Flores EM, Engelman EM. 1976. Apuntes sobre anatomía y morfología de las semillas de cactáceas. II. Caracteres de valor taxonómico. *Revista de Biología Tropical* **24**: 199-227.
- Flores J, Arredondo A, Jurado E. 2005. Comparative Seed Germination in Species of *Turbinicarpus*: an Endangered Cacti Genus. *Natural Areas Journal* **25**: 183-187.
- Flores J, Briones O. 2001. Plant life-form and germination in a Mexican inter-tropical desert: effects of soil water potential and temperature. *Journal of Arid Environments* **47**: 485-497. DOI: <https://doi.org/10.1006/jare.2000.0728>
- Flores J, Jurado E, Arredondo A. 2006. Effect of light on germination of seeds of Cactaceae from the Chihuahuan Desert, Mexico. *Seed Science Research* **16**: 149-155. DOI: <https://doi.org/10.1079/SSR2006242>
- Flores J, Jurado E, Chapa-Vargas L, Ceroni-Stuva A, Dávila-Aranda P, Galíndez G, Gurvich D, León-Lobos P, Ordóñez C, Ortega-Baes P, Ramírez-Bullón N, Sandoval A, Seal CE, Ullian T, Pritchard HW. 2011. Seeds photoblastism and its relationship with some plant traits in 136 cacti taxa. *Environmental and Experimental Botany* **71**: 79-88. DOI: <https://doi.org/10.1016/j.envexpbot.2010.10.025>
- Flores J, Jurado E, Jiménez-Bremont JF. 2008. Breaking seed dormancy in specially protected *Turbinicarpus lophophoroides* and *Turbinicarpus pseudopectinatus* (Cactaceae). *Plant Species Biology* **23**: 43-46. DOI: <https://doi.org/10.1111/j.1442-1984.2008.00206.x>

- Flores J, Pérez-Sánchez RM, Jurado E. 2017. The combined effect of water stress and temperature on seed germination of Chihuahuan Desert species. *Journal of Arid Environments* **146**: 95-98. DOI: <http://dx.doi.org/10.1016/j.jaridenv.2017.07.009>
- Fonseca RBS, Funch LS, Borba EL. 2012. Dispersão de sementes de *Melocactus glaucescens* e *M. paucispinus* (Cactaceae), no Município de Morro do Chapéu, Chapada Diamantina - BA. *Acta Botanica Brasileira* **26**: 481-492. DOI: <http://dx.doi.org/10.1590/S0102-33062012000200024>
- Font Quer P. 1973. *Diccionario de Botánica*. Barcelona: Editorial Labor S. A. ISBN: 84-335-5804-8
- Franco-Estrada D, Dávila-Aranda PD, Téllez-Valdés O. 2014. *Las cactáceas columnares del Valle de Tehuacán-Cuicatlán*. México DF: Facultad de Estudios Superiores Iztacala, Universidad Nacional Autónoma de México.
- Funes G, Díaz S, Venier P. 2009. La temperatura como principal determinante de la germinación en especies del Chaco seco de Argentina. *Ecología Austral* **19**: 129-138.
- García-Beltrán JÁ, Barrios D, Cuza-Pérez A. 2017. Heteromorphism in seeds of *Leptocereus scopulophilus* (Cactaceae) from Pan de Matanzas, Cuba. *Seed Science Research* **27**: 311-320. DOI: <https://doi.org/10.1017/S0960258517000289>
- Godínez-Álvarez H, Valiente-Banuet A. 1998. Germination and early seedling growth of Tehuacan Valley cacti species: the role of soils and seed ingestion by dispersers on seedling growth. *Journal of Arid Environments* **39**: 21-31. DOI: <https://doi.org/10.1006/jare.1998.0376>
- Godínez-Álvarez H, Valiente-Banuet A. 2000. Fruit-feeding behavior of the bats *Leptonycteris curasoae* and *Choeronycteris mexicana* in flight cage experiments: consequences for dispersal of columnar cactus Seeds. *Biotropica* **32**: 552-556. DOI: [http://dx.doi.org/10.1646/0006-3606\(2000\)032\[0552:FFBOTB\]2.0.CO;2](http://dx.doi.org/10.1646/0006-3606(2000)032[0552:FFBOTB]2.0.CO;2)
- Godínez-Álvarez H, Valiente-Banuet A, Rojas-Martínez A. 2002. The role of seed dispersers in the population dynamics of the columnar Cactus *Neobuxbaumia tetetzo*. *Ecology* **83**: 2617-2629. DOI: [http://dx.doi.org/10.1890/0012-9658\(2002\)083\[2617:TROSDI\]2.0.CO;2](http://dx.doi.org/10.1890/0012-9658(2002)083[2617:TROSDI]2.0.CO;2)
- Godínez-Álvarez H, Valverde T, Ortigas-Baes P. 2003. Demographic trends in Cactaceae. *The Botanical Review* **69**: 173-203. DOI: [http://dx.doi.org/10.1663/0006-8101\(2003\)069\[0173:DTITC\]2.0.CO;2](http://dx.doi.org/10.1663/0006-8101(2003)069[0173:DTITC]2.0.CO;2)
- Goettsch B, Hilton-Taylor C, Cruz-Pinon G, Duffy JP, Frances A, Hernandez H, Inger R, Pollock C, Schipper J, Superina M, Taylor NP, Tognelli M, Abba AM, Arias S, Arreola-Nava HJ, Baker MA, Bárcenas RT, Barrios D, Braun P, Butterworth CA, Búrquez, A, Cáceres F, Chazaro-Basañez, M, Corral-Díaz R, Perea M del V, Demaio PH, de Barros, WAD, Durán, R, Yancas LF, Felger RS, Fitz-Maurice, B., Fitz-Maurice WA, Gann G, Gómez-Hinostrosa C, Gonzales-Torres, LR, Patrick Griffith M, Guerrero PC, Hammel B, Heil KD, Hernández-Oria JG, Hoffmann M, Ishihara MI, Kiesling R, Larocca J, León-de la Luz JL, Loaiza CR, Lowry M, Machado MC, Majure LC, Martínez-Ávalos JG, Martorell C, Maschinski J, Méndez E, Mittermeier RA, Nassar JM, Negrón-Ortiz V, Oakley LJ, Ortega-Baes P, Ferreira ABP, Pinkava DJ, Porter JM, Puente-Martínez R, Gamarra JR, Saldivia-Pérez P, Sánchez-Martínez E, Smith M, Sotomayor JM, Stuart SN, Tapia-Muñoz JL, Terrazas T, Terry M, Trevisson M, Valverde T, Devender TRV, Véliz-Pérez ME, Walter HE, Wyatt SA, Zappi D, Zavala-Hurtado JA, Gaston KJ. 2015. High proportion of cactus species threatened with extinction. *Nature Plants* **1**: 15142. DOI: <https://doi.org/10.1038/NPLANTS.2015.142>
- Gomes VGN, Quirino ZGM, Araujo HFP. 2014. Frugivory and seed dispersal by birds in *Cereus jamacaru* DC. ssp. *jamacaru* (Cactaceae) in the Caatinga of Northeastern Brazil. *Brazilian Journal of Biology* **74**: 32-40. DOI: <http://dx.doi.org/10.1590/1519-6984.15312>
- Gonzalez-Cortés A, Reyes-Valdés MH, Robledo-Torres V, Villarreal-Quintanilla JA, Ramírez-Godina F. 2018. Pre-germination treatments in four prickly pear cactus (*Opuntia* sp.) species from Northeastern Mexico. *Australian Journal of Crop Science* **12**: 1676-1684. DOI: <https://doi.org/10.21475/ajcs.18.12.10.pne1430>
- Goodman J, Walters D, Bradley K, Maschinski J, Salazar A. 2012. Seeds of endangered *Harrisia fragrans* form persistent soil seed banks and withstand orthodox storage conditions. *Haseltonia* **18**: 85-94. DOI: <http://dx.doi.org/10.2985/026.018.0111>
- Gorai M, El Aloui W, Yang X, Neffati M. 2014. Toward understanding the ecological role of mucilage in seed germination of a desert shrub *Henophyton deserti*: interactive effects of temperature, salinity and osmotic stress. *Plant and Soil* **374**: 727-738. DOI: <https://doi.org/10.1007/s11104-013-1920-9>
- Griffith MP. 2004. Early cactus evolution: the postmodern view. *Haseltonia* **10**: 3-11.
- Guerrero PC, Mardones D, Viveros N, Peña-Gómez FT, Bustamante RO. 2016. Evolutionary change in the germination niche between related species within *Neoporteria* clade (Cactaceae) is idiosyncratic to habitat type. *Gayana Botanica* **73**: 177-182. DOI: <https://doi.org/10.4067/S0717-66432016000200177>
- Guillén S, Benítez J, Martínez-Ramos M, Casas A. 2009. Seed germination of wild, in situ-managed, and cultivated populations of columnar cacti in the Tehuacán-Cuicatlán Valley, Mexico. *Journal of Arid Environments* **73**: 407-413. DOI: <https://doi.org/10.1016/j.jaridenv.2008.12.018>

- Guillén S, Casas A, Rodríguez-Morales J. 2015. Patrones de germinación asociado a la domesticación y a la historia natural de cactáceas columnares del Valle de Tehuacán-Cuicatlán, México. *Gaia Scientia* **9**: 117-128.
- Guillén S, Terrazas T, De la Barrera E, Casas A. 2011. Germination differentiation patterns of wild and domesticated columnar cacti in a gradient of artificial selection intensity. *Genetic Resources and Crop Evolution* **58**: 409-423. DOI: <https://doi.org/10.1007/s10722-010-9586-0>
- Günster A. 1992. Aerial seed banks in the Central Namib: distribution of serotinous plants in relation to climate and habitat. *Journal of Biogeography* **19**: 563-572. DOI: <https://doi.org/10.2307/2845775>
- Gurvich DE, Funes G, Giorgis MA, Demaio P. 2008. Germination characteristics of four Argentinean endemic *Gymnocalycium* (Cactaceae) species with different flowering phenologies. *Natural Areas Journal* **28**: 104-108. DOI: [http://dx.doi.org/10.3375/0885-8608\(2008\)28\[104:GCOFAE\]2.0.CO;2](http://dx.doi.org/10.3375/0885-8608(2008)28[104:GCOFAE]2.0.CO;2)
- Gurvich DE, Pérez-Sánchez R, Bauk K, Jurado E, Ferrero MC, Funes G, Flores J. 2017. Combined effect of water potential and temperature on seed germination and seedling development of cacti from a mesic Argentine ecosystem. *Flora* **227**: 18-24. DOI: <https://doi.org/10.1016/j.flora.2016.12.003>
- Haddaway NR, Collins AM, Coughlin D, Kirk S. 2015. The role of Google Scholar in evidence reviews and its applicability to grey literature searching. *PLoS ONE* **10**: e0138237. DOI: <https://dx.doi.org/10.1371/journal.pone.0138237>
- Hernández-Ledesma P, Berendsohn WG, Borsch T, Von Mering S, Akhani H, Arias S, Castañeda-Noa I, Eggli U, Eriksson R, Flores-Olvera H, Fuentes-Bazán S, Kadereit G, Klak C, Korotkova N, Nyffeler R, Ocampo G, Ochoterena H, Oxelman B, Rabeler RK, Sanchez A, Schlumberger BO, Uotila P. 2015. A taxonomic backbone for the global synthesis of species diversity in the angiosperm order Caryophyllales. *Willdenowia* **45**: 281-383. DOI: <https://doi.org/10.3372/wi.45.45301>
- Hunt D, Taylor NP, Graham C. 2006. The New Cactus Lexicon. England: dh books. ISBN: 0953813444
- Jiménez-Sierra CL, Matías-Palafox ML. 2015. Dinámica poblacional de cactáceas amenazadas. Universidad Autónoma Metropolitana, Unidad Iztapalapa. Informe final SNIB-CONABIO proyecto No. HK026. <http://www.conabio.gob.mx/institucion/proyectos/resultados/InfHK026.pdf> (accessed March 16, 2020)
- Jurado E, Flores J. 2005. Is seed dormancy under environmental control or bound to plant traits? *Journal of Vegetation Science* **16**: 559-564. DOI: <https://doi.org/10.1111/j.1654-1103.2005.tb02396.x>
- Kin AG, Mazzola MB, Cenizo VJ. 2015. Seed germination and seedling growth of the geophytic *Pterocactus tuberosus* (Cactaceae). *Journal of the Torrey Botanical Society* **142**: 283-291. DOI: <https://doi.org/10.3159/TORREY-D-14-00056.1>
- Lasso E, Barrientos LS. 2015. Epizooticoria por medio de iguanas en el bosque seco: ¿Un mecanismo de dispersión de semillas pasado por alto? *Colombia Forestal* **18**: 151-159. DOI: <http://dx.doi.org/10.14483/udistrital.jour.colomb.for.2015.1.a09>
- Lenzi M, Graipel ME, Matos JZd, Fraga AM, Orth AI. 2012. Dispersão zoocórica e hidroocórica marítima de *Opuntia monacantha* (Willd.) Haw. (Cactaceae). *Biotemas* **25**: 47-53. DOI: <https://doi.org/10.5007/2175-7925.2012v25n1p47>
- Leuenberger BE. 1986. *Pereskia* (Cactaceae). *Memoirs of the New York Botanical Garden* **41**: 1-141. ISBN 10: 0-89327-307-4; ISBN-13: 978-0893273071
- Lima AT, Meiado MV. 2017. Discontinuous hydration alters seed germination under stress of two populations of cactus that occur in different ecosystems in Northeast Brazil. *Seed Science Research* **27**: 292-302. DOI: <https://doi.org/10.1017/S0960258517000241>
- Lima AT, Meiado MV. 2018. Effects of seed hydration memory on initial growth under water deficit of cactus from two populations that occur in different ecosystems in Northeast Brazil. *Plant Species Biology* **33**: 268-275. DOI: <https://doi.org/10.1111/1442-1984.12219>
- Lindow-López L, Galíndez G, Aparicio-González M, Sühling S, Rojas-Aréchiga M, Pritchard HW, Ortega-Baes P. 2018b. Effects of alternating temperature on cactus seeds with a positive photoblastic response. *Journal of Arid Environments* **148**: 74-77. DOI: <http://dx.doi.org/10.1016/j.jaridenv.2017.10.006>
- Lindow-López L, Galíndez G, Sühling S, Pastrana-Ignes V, Gorostiague P, Gutiérrez A, Ortega-Baes P. 2018a. Do cacti form soil seed banks? An evaluation using species from the Southern Central Andes. *Plant Biology* **20**: 1053-1058. DOI: <https://doi.org/10.1111/plb.12868>
- Lone AB, Colombo RC, Andrade BLG, Takahashia LSA, Faria RT. 2016. Physical characterization of *Rhipsalis* (Cactaceae) fruits and seeds germination in different temperatures and light regimes. *Brazilian Journal of Biology* **76**: 367-373. DOI: <http://dx.doi.org/10.1590/1519-6984.15914>
- López-Escamilla AL, López-Herrera M, Loaiza-Alanis C. 2016. Effect of different gelling agents on *in vitro* germination and seedling development of *Echinocactus platyacanthus* Link et Otto (Cactaceae). *Polibotánica* **42**: 153-166. DOI: <https://doi.org/10.18387/polibotanica.42.8>
- López-Gómez R, Sánchez-Romero P. 1989. Germinación de dos variedades de pitaya *Stenocereus griseus*

- (Haworth) Buxbaum. *Cactáceas y Suculentas Mexicanas* **34**: 35-40.
- López-Urrutia E, Martínez-García M, Monsalvo-Reyes A, Salazar-Rojas V, Montoya R, Campos JE. 2014. Differential RNA-and protein-expression profiles of cactus seeds capable of hydration memory. *Seed Science Research* **24**: 91-99. DOI: <https://doi.org/10.1017/S0960258513000317>
- Loustalot E, Malda GX, Suzán H, Hernández LG, Guevara A. 2014. Estudio de germinación y crecimiento en semillas de *Ferocactus histrix* (De Candolle). *Cactáceas y Succulentas Mexicanas* **59**: 79-95.
- Loza-Cornejo S, Terrazas T, López-Mata L. 2012. Fruits, seeds and germination in five species of globose Cactaceae (Cactaceae). *Interciencia* **37**: 197-203.
- Luna B, Chamorro D. 2016. Germination sensitivity to water stress of eight Cistaceae species from the western mediterranean. *Seed Science Research* **26**: 101-110. DOI: <https://doi.org/10.1017/S096025851600009X>
- Maiti RK, Hernández-Piñero JL, Valdez-Marroquín M. 1994. Seed ultrastructure and germination of some species of Cactaceae. *Phyton* **55**: 97-105.
- Maiti RK, Singh VP, Baquie A, Sánchez-Arreola E, Wesche-Ebeling P, Cuervo-Parra JA, Perdome-Velázquez H, Lorenzo JL. 2003. *Cactus: Biology, Propagation and Conservation*. Chicago: Gurav Society of Agricultural Research Information Centre.
- Maldonado C, Pujado E, Squeo FA. 2002. El efecto de la disponibilidad de agua durante el crecimiento de *Lycopersicon chilense* sobre la capacidad de sus semillas para germinar a distintas temperaturas y concentraciones de manitol y NaCl. *Revista Chilena de Historia Natural* **75**: 651-660. DOI: <http://dx.doi.org/10.4067/S0716-078X2002000400002>
- Mandujano MC, Golubov J, Montaña C. 1997. Dormancy and endozoochorous dispersal of *Opuntia rastrera* seeds in the southern Chihuahuan Desert. *Journal of Arid Environments* **36**: 259-266. DOI: <https://doi.org/10.1006/jare.1996.0210>
- Mandujano MC, Montaña C, Rojas-Aréchiga M. 2005. Breaking seed dormancy in *Opuntia rastrera* from the Chihuahuan desert. *Journal of Arid Environments* **62**: 15-21. DOI: <https://doi.org/10.1016/j.jaridenv.2004.10.009>
- Mandujano MC, Golubov J, Rojas-Aréchiga M. 2007. Efecto del ácido giberélico en la germinación de tres especies del género *Opuntia* (Cactaceae) del Desierto Chihuahuense. *Cactáceas y Suculentas Mexicanas* **52**: 46-52.
- Marchi MNG, Civatti LM, Viana CM, de Assis JGA, Bellintani MC, de Santana JRF. 2013. Seed cryopreservation of the native cacti *Discocactus zehntneri*, *Pilosocereus gounellei* and *Stephanocereus luetzelburgii* from Bahia, Brazil. *African Journal of Biotechnology* **12**: 3250-3254. DOI: <https://doi.org/10.5897/AJB2012.10391>
- Martínez-Berdeja A, Ezcurra E, Sanders AC. 2015. Delayed seed dispersal in California deserts. *Madroño* **62**: 21-32. DOI: <https://doi.org/10.3120/0024-9637-62.1.21>
- Martins LST, Pereira TS, Carvalho ASDR, Barros C, De Andrade ACS. 2012. Seed germination of *Pilosocereus arrabidae* (Cactaceae) from a semiarid region of south-east Brazil. *Plant Species Biology* **27**: 191-200. DOI: <https://doi.org/10.1111/j.1442-1984.2011.00360.x>
- Mascot-Gómez E, Flores J, López-Lozano NE, Yáñez-Espinosa L. 2020. Seed germination of Southern Chihuahuan desert cacti: Effect of mucilage, light and phytohormones. *Flora* **263**: 151528. DOI: <https://doi.org/10.1016/j.flora.2019.151528>
- Matías-Palafox ML. 2007. *Estructura poblacional y biología reproductiva de Turbinicarpus horripilus (Lem.) Vác. John y Riha (Cactaceae)*. MS Thesis, Universidad Autónoma Metropolitana.
- Matías-Palafox ML, Jiménez-Sierra C. 2006. Evaluación del riesgo de extinción de las poblaciones de *Turbinicarpus horripilus* (Lem.) Vác. John & Riha, cactácea endémica de la Barranca de Metztitlán, México. *Cactáceas y Succulentas Mexicanas* **51**: 100-110.
- Mazzola MB, Cenizo VJ, Kin AG. 2013. Factores que afectan la germinación de *Trichocereus candicans* (Cactaceae). *Boletín de la Sociedad Argentina de Botánica* **48**: 515-523.
- McDonough WT. 1964. Germination responses of *Carnegiea gigantea* and *Lemaireocereus thurberi*. *Ecology* **45**: 155-159. DOI: <https://doi.org/10.2307/1937116>
- Meiádo MV, Correa LS, Rocha EA, Rojas-Aréchiga M, Leal IR. 2010. Seed germination responses of *Cereus jamacaru* DC. ssp. *jamacaru* (Cactaceae) to environmental factors. *Plant Species Biology* **25**: 120-128. DOI: <https://doi.org/10.1111/j.1442-1984.2010.00274.x>
- Meiádo MV, Rojas-Aréchiga M, Siqueira-Filhos JA, Leal IR. 2016. Effects of light and temperature on seed germination of cacti of Brazilian ecosystems. *Plant Species Biology* **31**: 87-97. DOI: <https://doi.org/10.1111/1442-1984.12087>
- Milberg P, Andersson L, Thompson K. 2000. Large-seeded species are less dependent on light for germination than small-seeded ones. *Seed Science Research* **10**: 99-104. DOI: <https://doi.org/10.1017/S0960258500000118>
- Montiel S, Montaña C. 2000. Vertebrate frugivory and seed dispersal of a Chihuahuan Desert cactus. *Plant Ecology* **146**: 221-229. DOI: <https://doi.org/10.1023/A:1009819419498>

- Montiel S, Montaña C. 2003. Seed bank dynamics of the desert cactus *Opuntia rastrera* in two habitats from the Chihuahuan Desert. *Plant Ecology* **166**: 241-248. DOI: <https://doi.org/10.1023/A:1023255314277>
- Munguía-Rosas MA, Jácome-Flores ME, Sosa VJ, Quiroz-Cerón LM. 2009. Removal of *Pilosocereus leucocephalus* (Cactaceae, tribe Cereeae) seeds by ants and their potential role as primary seed dispersers. *Journal of Arid Environments* **73**: 578-581. DOI: <https://doi.org/10.1016/j.jaridenv.2008.12.017>
- Naranjo ME, Rengifo C, Soriano PJ. 2003. Effect of ingestion by bats and birds on seed germination of *Stenocereus griseus* and *Subpilocereus repandus* (Cactaceae). *Journal of Tropical Ecology* **19**: 19-25. DOI: <https://doi.org/10.1017/S0266467403003031>
- Nascimento JPB, Meiado MV, Nicola PA, Pereira LCM. 2015. Germinação de sementes de *Tacinga inamoena* (K. Schum.) N.P. Taylor & Stuppy (Cactaceae) após endozoocoria por *Chelonoidis carbonaria* (Spix, 1824) (Reptilia: Testudinidae). *Gaia Scientia* **9**: 9-14.
- Nolasco H, Vega-Villasante F, Diaz-Rondero A. 1997. Seed germination of *Stenocereus thurberi* (Cactaceae) under different solar irradiation levels. *Journal of Arid Environments* **36**: 123-132. DOI: <https://doi.org/10.1006/jare.1996.0199>
- Nolasco H, Vega-Villasante F, Romero-Schmidt HL, Diaz-Rondero A. 1996. The effects of salinity, acidity, light and temperature on the germination of seeds of cardón (*Pachycereus pringlei* (S. Wats.) Britton & Rose, Cactaceae). *Journal of Arid Environments* **33**: 87-94. DOI: <https://doi.org/10.1006/jare.1996.0048>
- Oliveira DB, Lima AT, Rocha EA, Meiado MV. 2017. O aumento da temperatura reduz a tolerância ao estresse hídrico na germinação de sementes de *Pereskia grandifolia* Haw. subsp. *grandifolia* (Cactaceae)? *Gaia Scientia* **11**: 26-36. DOI: <https://doi.org/10.22478/ufpb.1981-1268.2017v11n4.35466>
- Olvera-Carrillo Y, Márquez-Guzmán J, Barradas VL, Sánchez-Coronado ME, Orozco-Segovia A. 2003. Germination of the hard seed coated *Opuntia tomentosa* S.D., a cacti from the México valley. *Journal of Arid Environments* **55**: 29-42. DOI: [https://doi.org/10.1016/S0140-1963\(02\)00268-9](https://doi.org/10.1016/S0140-1963(02)00268-9)
- Olvera-Carrillo Y, Márquez-Guzmán J, Sánchez-Coronado ME, Barradas VL, Rincón E, Orozco-Segovia A. 2009. Effect of burial on the germination of *Opuntia tomentosa*'s (Cactaceae, Opuntioideae) seeds. *Journal of Arid Environments* **73**: 421-427. DOI: <https://doi.org/10.1016/j.jaridenv.2008.12.011>
- Ordoñez-Salanueva CA, Orozco-Segovia A, Canales-Martínez M, Seal CE, Pritchard HW, Flores-Ortiz CM. 2017. Ecological longevity of *Polaskia chende* (Cactaceae) seeds in the soil seed bank, seedling emergence and survival. *Plant Biology* **19**: 973-982. DOI: <https://doi.org/10.1111/plb.12611>
- Ordoñez-Salanueva CA, Seal CE, Pritchard HW, Orozco-Segovia A, Canales-Martínez M, Flores-Ortiz CM. 2015. Cardinal temperatures and thermal time in *Polaskia Backeb* (Cactaceae) species: Effect of projected soil temperature increase and nurse interaction on germination timing. *Journal of Arid Environments* **115**: 73-80. DOI: <http://dx.doi.org/10.1016/j.jaridenv.2015.01.006>
- Orozco-Segovia A, Márquez-Guzmán J, Sánchez-Coronado ME, Gamboa-de Buen A, Baskin JM, Baskin CC. 2007. Seed Anatomy and Water Uptake in Relation to Seed Dormancy in *Opuntia tomentosa* (Cactaceae, Opuntioideae). *Annals of Botany* **99**: 581-592. DOI: <https://doi.org/10.1093/aob/mcm001>
- Ortega-Baes P, Aparicio-González M, Galíndez G, del Fueyo P, Sühling S, Rojas-Aréchiga M. 2010. Are cactus growth forms related to germination responses to light? A test using *Echinopsis* species. *Acta Oecologica* **36**: 339-342. DOI: <https://doi.org/10.1016/j.actao.2010.02.006>
- Ortega-Baes P, Galíndez G, Sühling S, Rojas-Aréchiga M, Daws MI, Pritchard HW. 2011. Seed germination of *Echinopsis schickendantzii* (Cactaceae): the effects of constant and alternating temperatures. *Seed Science & Technology* **39**: 219-224. DOI: <https://doi.org/10.15258/sst.2011.39.1.21>
- Ortega-Baes P, Rojas-Aréchiga M. 2007. Seed germination of *Trichocereus terscheckii* (Cactaceae): Light, temperature and gibberellic acid effects. *Journal of Arid Environments* **69**: 169-176. DOI: <https://doi.org/10.1016/j.jaridenv.2006.09.009>
- Ortiz TA, Gomes GR, Takahashi LSA, Urbano MR, Strapasson E. 2014. Water and salt stress in germinating seeds of pitaya genotypes (*Hylocereus* spp.). *African Journal of Agricultural Research* **9**: 3610-3619. DOI: <https://doi.org/10.5897/AJAR2014.9114>
- Patané C, Saita A, Tubeileh A, Cosentino SL, Cavallaro V. 2016. Modeling seed germination of unprimed and primed seeds of sweet sorghum under PEG-induced water stress through the hydrotimic analysis. *Acta Physiologiae Plantarum* **38**: 115-127. DOI: <https://doi.org/10.1007/s11738-016-2135-5>
- Pérez-Sánchez RM, Jurado E, Chapa-Vargas L, Flores J. 2011. Seed germination of Southern Chihuahuan Desert plants in response to elevated temperatures. *Journal of Arid Environments* **75**: 978-980. DOI: <https://doi.org/10.1016/j.jaridenv.2011.04.020>
- Pérez-Villafañá MG, Valiente-Banuet A. 2009. Effectiveness of dispersal of an ornithocorous cactus *Myrtillocactus geometrizans* (Cactaceae) in a patchy environment. *Open Biology Journal* **2**: 101-113. DOI: <https://doi.org/10.2174/1874196700902010101>

- Peters EM, Martorell C, Ezcurra E. 2009. The adaptive value of cued seed dispersal in desert plants: Seed retention and release in *Mammillaria pectinifera* (Cactaceae), a small globose cactus. *American Journal of Botany* **96**: 537-541. DOI: <https://doi.org/10.3732/ajb.0800157>
- Podda L, Santo A, Leone C, Mayoral O, Bacchetta G. 2017. Seed germination, salt stress tolerance and seedling growth of *Opuntia ficus-indica* (Cactaceae), invasive species in the Mediterranean Basin. *Flora* **229**: 50-57. DOI: <https://doi.org/10.1016/j.flora.2017.02.002>
- Potter RL, Petersen JL, Ueckert DN. 1984. Germination responses of *Opuntia* spp. to temperature, scarification, and other seed treatments. *Weed Science Society of America* **32**: 106-110. DOI: <https://doi.org/10.1017/S0043174500058598>
- Ramírez-González G, Rodríguez-De la O JL, Martínez-Solís J, Colinas-León MT. 2019. Germinación y crecimiento ex vitro e in vitro de cinco especies de cactáceas del género *Mammillaria*. *Polibotánica* **48**: 99-110. DOI: <https://doi.org/10.18387/polibotanica.48.8>
- Ramírez-Padilla CA, Valverde T. 2005. Germination responses of three congeneric cactus species (*Neobuxbaumia*) with differing degrees of rarity. *Journal of Arid Environments* **61**: 333-343. DOI: <https://doi.org/10.1016/j.jaridenv.2004.09.006>
- Rito KF, Rocha EA, Leal IR, Meiado MV. 2009. As sementes de mandacaru têm memória hídrica? *Boletín de la Sociedad Latinoamericana y del Caribe de Cactáceas y otras Suculentas* **6**: 26-31.
- Rodríguez-Morales J, Guillen S, Casas A. 2013. Consecuencias de la domesticación de *Stenocereus stellatus* en el tamaño de las semillas y en la germinación en un gradiente de estrés hídrico. *Botanical Sciences* **91**: 485-492. DOI: <https://doi.org/10.17129/botsci.425>
- Rodríguez-Ortega C, Franco M, Mandujano MC. 2006. Serotiny and seed germination in three threatened species of *Mammillaria* (Cactaceae). *Basic and Applied Ecology* **7**: 533-544. DOI: <https://doi.org/10.1016/j.baae.2006.04.001>
- Rodríguez-Ruiz ER, Poot-Poot WA, Rangel-Lucio JA, Vaquera-Huerta H, González-Gaona OJ, Treviño-Carreón J. 2018. Germinación in vitro de biznaga cabuchera. *Revista Mexicana de Ciencias Agrícolas* **9**: 691-699.
- Rojas-Aréchiga M. 2008. Efecto del ácido giberélico en la germinación de cuatro especies del género *Mammillaria* del Valle de Tehuacán-Cuicatlán, México. *Boletín de la Sociedad Latinoamericana y del Caribe de Cactáceas y Suculentas* **5**: 21-23.
- Rojas-Aréchiga M. 2012. La importancia de la semilla en Cactaceae para estudios taxonómicos y filogenéticos. *Boletín de la Sociedad Latinoamericana y del Caribe de Cactáceas y otras Suculentas* **9**: 15-18.
- Rojas-Aréchiga M, Aguilar KM, Golubov J, Mandujano MC. 2011. Effect of gibberellic acid on germination of seeds of five species of cacti from the Chihuahuan desert, northern Mexico. *The Southwestern Naturalist* **56**: 393-400. DOI: <https://doi.org/10.1894/N01-DW-126.1>
- Rojas-Aréchiga M, Batis AI. 2001. Las semillas de cactáceas...¿forman bancos en el suelo? *Cactáceas y Suculentas Mexicanas* **46**: 76-82.
- Rojas-Aréchiga M, Casas A, Vázquez-Yanes CJ. 2001. Seed germination of wild and cultivated *Stenocereus stellatus* (Cactaceae) from the Tehuacán-Cuicatlán Valley, Central México. *Journal of Arid Environments*, **49**: 279-287. DOI: <https://doi.org/10.1006/jare.2001.0789>
- Rojas-Aréchiga M, Mandujano-Sánchez MC. 2009. Nuevo registro de semillas vivíparas en dos especies de cactáceas. *Cactáceas y Suculentas Mexicanas* **54**: 123-127.
- Rojas-Aréchiga M, Mandujano-Sánchez MC. 2017. Latencia secundaria en especies de la tribu Cactaeae (Cactaceae). *Polibotánica* **44**: 137-145. DOI: <https://doi.org/10.18387/polibotanica.44.10>
- Rojas-Aréchiga M, Mandujano MC, Golubov JK. 2013. Seed size and photoblastism in species belonging to tribe Cactaeae (Cactaceae). *Journal of Plant Research* **126**: 373-386. DOI: <https://doi.org/10.1007/s10265-012-0526-2>
- Rojas-Aréchiga M, Orozco-Segovia A, Vázquez-Yanes C. 1997. Effect of light on germination of seven species of cacti from the Zapotitlán Valley in Puebla, México. *Journal of Arid Environments* **36**: 571-578. DOI: <https://doi.org/10.1006/jare.1996.0218>
- Rojas-Aréchiga M, Vázquez-Yanes C. 2000. Cactus seed germination: a review. *Journal of Arid Environments* **44**: 85-104. DOI: <https://doi.org/10.1006/jare.1999.0582>
- Rojas-Aréchiga M, Vázquez-Yanes C, Orozco-Segovia A. 1998. Seed response to temperature of Mexican cacti species from two life forms: an ecophysiological interpretation. *Plant Ecology* **135**: 207-214. DOI: <https://doi.org/10.1023/A:1009757227493>
- Romero-Méndez U, Becerra-López JL, García-De la Peña C, Martínez-Ríos JJ, Czaja A, Muro G, Sánchez JA. 2018. Hydrochory in *Astrophytum coahuilense*: experiment to identify seminal structures that provide its buoyancy. *Polibotánica* **46**: 149-157. DOI: <https://doi.org/10.18387/polibotanica.46.8>
- Romero-Schmidt HL, Vega-Villasante F, Nolasco H, Montaña C. 1992. The effect of darkness, freezing, acidity and salinity on seed germination of *Ferocactus peninsulae* (Cactaceae). *Journal of Arid Environments* **23**: 389-395. DOI: [https://doi.org/10.1016/S0140-1963\(18\)30612-8](https://doi.org/10.1016/S0140-1963(18)30612-8)

- Romo-Campos L, Flores-Flores JL, Flores J, Álvarez-Fuentes G. 2010. Seed germination of *Opuntia* species from an aridity gradient in Central Mexico. *Journal of the Professional Association for Cactus Development* **12**: 181-198.
- Rosas-López U, Collazo-Ortega M. 2004. Conditions for the germination and the early growth of seedlings of *Polaskia chichipe* (Goss.) Backeberg and *Echinocactus platyacanthus* Link and Otto fa. *grandis* (Rose) Bravo-Hollis (Cactaceae). *International Journal of Experimental Botany* **53**: 213-220.
- Royal Botanic Gardens Kew. 2019. Seed Information Database (SID). v.7.1. <http://data.kew.org/sid/> (accessed August 20, 2019).
- Ruedas M, Valverde T, Castillo-Agüero S. 2000. Respuesta germinativa y crecimiento de plántulas de *Mammillaria magnimamma* (Cactaceae) bajo diferentes condiciones ambientales. *Boletín de la Sociedad Botánica de México* **66**: 25-35. DOI: <https://doi.org/10.17129/botsci.1608>
- Ruiz-González SP, Rojas-Aréchiga M, Mandujano MC. 2011. Descripción morfológica y germinación de las semillas de *Echinomastus unguispinus*. *Cactáceas y Succulentas Mexicanas* **56**: 36-44.
- Salas-Cruz LR, Foroughbackch-Pournabav R, Díaz-Jiménez ML, Cárdenas-Ávila ML, Flores-Valdes A. 2011. Germinación in vitro de cactáceas, utilizando zeolita como sustrato alternativo. *Revista Mexicana de Ciencias Agrícolas Publicación Especial*: 565-575.
- Salazar A, Maschinski J, Powell D. 2013. *Ex-Situ* Seed Conservation of Endangered Key Tree Cactus (*Pilosocereus robinii*). *Journal of Biodiversity & Endangered Species* **1**: 111-114. DOI: <https://doi-org/10.4172/jbes.1000111>
- Sánchez JA, Furrázola E, Pernús M, Torres-Arias Y. 2018. Síntesis y perspectivas. In: Sánchez JA, Furrázola E. eds. *Ecotecnologías para la restauración ecológica: los tratamientos de semillas y micorrizas*. La Habana: Editorial Academia, 93-96. ISBN: 978-959-270-384-1, 978-959-270-385-8
- Sánchez JA, Montejo L, Gamboa A, Albert-Puentes D, Hernández F. 2015. Germinación y dormancia de arbustos y trepadoras del bosque siempreverde de la Sierra del Rosario, Cuba. *Pastos y Forrajes* **38**: 11-28.
- Sánchez JA, Pernús M. 2018. Tratamientos de semillas. In: Sánchez JA, Furrázola E. eds. *Ecotecnologías para la restauración ecológica: los tratamientos de semillas y micorrizas*. La Habana: Editorial Academia, 5-54. ISBN: 978-959-270-384-1, 978-959-270-385-8
- Sánchez-Salas J, Flores J, Martínez-García E. 2006. Efecto del tamaño de semilla en la germinación de *Astrophytum myriostigma* Lemaire (Cactaceae) especie amenazada de extinción. *Interciencia* **31**: 371-375.
- Sánchez-Salas J, Jurado E, Flores J, Estrada-Castillón E, Muro-Pérez G. 2012. Desert species adapted for dispersal and germination during floods: Experimental evidence in two *Astrophytum* species (Cactaceae). *Flora* **207**: 707-711. DOI: <https://doi.org/10.1016/j.flora.2012.08.002>
- Sánchez-Salas Jurado E, Flores J, Muro-Pérez G, Estrada-Castillón E. 2013. Adaptaciones inusuales en semillas y plántulas del desierto: Mecanismos para su germinación & dispersión. *Boletín de la Sociedad Latinoamericana y del Caribe de Cactáceas y Succulentas* **10**: 19-23.
- Sánchez-Soto B, García-Moya E, Terrazas T, Reyes-Olivas A. 2005. Efecto de la hidratación discontinua sobre la germinación de tres cactáceas del desierto costero de Topolobampo, Ahome, Sinaloa. *Cactáceas y Succulentas Mexicanas* **50**: 4-14.
- Sánchez-Soto B, Reyes-Olivas Á, García-Moya E, Terrazas T. 2010. Germinación de tres cactáceas que habitan la región costera del noroeste de México. *Interciencia* **35**: 299-305.
- Sánchez-Venegas G. 1997. Germinación, viabilidad y características distintivas de la semilla de *Opuntia joconostle* Weber, forma cuaresmero. *Cactáceas y Succulentas Mexicanas* **42**: 16-21.
- Santini BA, Martorell C. 2013. Does retained-seed priming drive the evolution of serotiny in drylands? An assessment using the cactus *Mammillaria hernandezii*. *American Journal of Botany* **100**: 365-373. DOI: <https://doi.org/10.3732/ajb.1200106>
- Santini BA, Rojas-Aréchiga M, García-Morales E. 2017. Priming effect on seed germination: Is it always positive for cacti species? *Journal of Arid Environments* **147**: 155-158. DOI: <http://dx.doi.org/10.1016/j.jaridenv.2017.07.013>
- Santos LDN, Pereira IMS, Ribeiro JR, Las-Casas FMG. 2019. Frugivoria por aves em quatro espécies de Cactaceae na Caatinga, uma floresta seca no Brasil. *Iheringia, Série Zoologia* **109**: e2019034. DOI: <https://doi.org/10.1590/1678-4766e2019034>
- Seal CE, Daws MI, Joel F, Ortega-Baes P, Galíndez G, León-Lobos P, Sandoval A, Stuva AC, Ramírez N, Dávila-Aranda P et al. 2017. Thermal buffering capacity of the germination phenotype across the environmental envelope of the Cactaceae. *Global Change Biology* **23**: 5309-5317. DOI: <https://doi.org/10.1111/gcb.13796>
- Seal CE, Flores J, Ceroni-Stuva A, Dávila-Aranda P, León-Lobos P, Ortega-Baes P, Galíndez G, Aparicio-González MA, Castro-Cepero V, Daws MI, Eason M, Flores-Ortiz CM, del Fueyo PA, Olwell P, Ordoñez C, Peñalosa-Castro I, Quintanar-Zúñiga R, Ramírez-Bullón N, Rojas-Aréchiga M, Rosas M, Sandoval A, Stuppy W, Ulian T, Vázquez-Medrano J, Walter H, Way M, Pritchard HW. 2009. The Cactus Seed Biology Database (release 7.1). Royal Botanic Gardens, Kew. <https://data.kew.org/sid/> (accessed August 29, 2020)

- Secorun AC, Souza LA. 2011. Morphology and anatomy of *Rhipsalis cereuscula*, *Rhipsalis floccosa* subsp. *hohenauensis* and *Lepismium cruciforme* (Cactaceae) seedlings. *Revista Mexicana de Biodiversidad* **82**: 131-143. DOI: <http://dx.doi.org/10.22201/ib.20078706e.2011.1.371>
- Silvius KM. 1995. Avian consumers of Cardón fruits (*Stenocereus griseus*: Cactaceae) on Margarita island, Venezuela. *Biotropica* **27**: 96-105. DOI: <https://dx.doi.org/10.2307/2388907>
- Simão E, Socolowski F, Takaki M. 2007. The epiphytic Cactaceae *Hylocereus setaceus* (Salm-Dick ex DC.) Ralf Bauer seed germination is controlled by light and temperature. *Brazilian archives of Biology and Technology* **40**: 655-662. DOI: <http://dx.doi.org/10.1590/S1516-89132007000400011>
- Simão E, Takaki M, Cardoso VJM. 2010. Germination response of *Hylocereus setaceus* (Salm-Dyck ex DC.) Ralf Bauer (Cactaceae) seeds to temperature and reduced water potentials. *Brazilian Journal of Biological Sciences* **70**: 135-144. DOI: <http://dx.doi.org/10.1590/S1519-69842010000100019>
- Sosa-Pivatto M, Funes G, Ferreras AE, Gurvich DE. 2014. Seed mass, germination and seedling traits for some central Argentinian cacti. *Seed Science Research* **24**: 71-77. DOI: <http://dx.doi.org/10.1017/S0960258513000366>
- Souza DFH, Marcos-Filho J. 2001. The seed coat as a modulator of see-environmental relationships in Fabaceae. *Revista Brasileira de Sementes* **4**: 365-375.
- Strittmatter LI, Negron-Ortiz V, Hickey RJ. 2002. Subdioecy in *Consolea spinosissima* (cactaceae): breeding system and embryological studies. *American Journal of Botany* **89**: 1373-1387. DOI: <https://doi.org/10.3732/ajb.89.9.1373>
- Thompson K, Grime JP. 1979. Seasonal variation in the seed banks of herbaceous species in ten contrasting habitats. *Journal Ecology* **67**: 893-921. DOI: <http://dx.doi.org/10.2307/2259220>
- Tribouillois H, Dürr C, Demilly D, Wagner M-H, Justes E. 2016. Determination of germination response to temperature and water potential for a wide range of cover crop species and related functional groups. *PLOS One* **11**: e0161185. DOI: <http://dx.doi.org/10.1371/journal.pone.0161185>
- Trujillo AG, Espinoza JL, Ortega R, Ávila NY, Espinosa AP. 2014. Efecto del tiempo de almacenamiento de la semilla en la germinación y sobrevivencia de *Ferocactus townsendianus* Britt & Rose. *Interciencia* **39**: 732-735.
- Valiente-Banuet A, Rojas-Martínez A, Casas A, Arizmendi MdC, Dávila P. 1997. Pollination biology of two winter-blooming giant columnar cacti in the Tehuacán Valley, central Mexico. *Journal of Arid Environments* **37**: 331-341. DOI: <https://doi.org/10.1006/jare.1997.0267>
- Valverde PL, Jimenez-Sierra C, Lopez-Ortega G, Zavala-Hurtado JA, Rivas-Arancibia S, Rendon-Aguilar B, Perez-Hernández MA, Cornejo-Romero A, Carrillo-Ruiz H. 2015. Floral morphometry, anthesis, and pollination success of *Mammillaria pectinifera* (Cactaceae), a rare and threatened endemic species of Central Mexico. *Journal of Arid Environments* **116**: 29-32. DOI: <http://dx.doi.org/10.1016/j.jaridenv.2015.01.016>
- Valverde PL, Zavala-Hurtado JA. 2006. Assessing the ecological status of *Mammillaria pectinifera* Weber (Cactaceae), a rare and threatened species endemic of the Tehuacán-Cuicatlán Region in Central Mexico. *Journal of Arid Environments* **64**: 193-208. DOI: <http://dx.doi.org/10.1016/j.jaridenv.2005.06.001>
- Vázquez-Castillo S, Miranda-Jácome A, Ruelas E. 2018. Patterns of frugivory in the columnar cactus *Pilosocereus leucocephalus*. *Ecology and Evolution* **9**: 268-1277. DOI: <http://dx.doi.org/10.1002/ece3.4833>
- Vega-Villasante F, Nolasco H, Montaña C, Romero-Schmidt H, Vega-Villasante E. 1996. Efecto de la temperatura, acidez, iluminación, salinidad, irradiación solar y humedad sobre la germinación de semillas de *Pachycereus pecten-aboriginum* “cardón barbón” (Cactaceae). *Cactáceas y Suculentas Mexicanas* **41**: 51-61.
- Veiga-Barbosa L, González-Benito ME, Assis JGA, Pérez-García F. 2010. Germination and cryopreservation of several cactus species from NE Brazil. *Seed Science & Technology* **38**: 218-224. DOI: <https://doi.org/10.15258/sst.2010.38.1.22>
- Venable DL. 2007. Bet hedging in a guild of desert annuals. *Ecology* **88**: 1086-1090.
- Venable DL, Lawlor L. 1980. Delayed germination and dispersal in desert annuals: escape in space and time. *Oecologia* **46**: 272-282. DOI: <https://dx.doi.org/10.1007/BF00540137>
- Villanueva RM, Navarro MdC, Eliosa HR. 2016. Germinación de tres especies de cactáceas endémicas de México en condiciones asépticas. *Zonas Áridas* **16**: 1-16. DOI: <http://dx.doi.org/10.21704/za.v16i1.633>
- Voit G. 1979. Untersuchungen zur Mikromorphologie der Cactaceen-Samen unter Berücksichtigung taxonomische Aspekte. PhD. Thesis, University of Heidelberg, Alemania.
- Walck JL, Baskin JM, Baskin CC, Hidayati SN. 2005. Defining transient and persistent seed banks in species with pronounced seasonal dormancy and germination patterns. *Seed Science Research* **15**: 189-196. DOI: <https://doi.org/10.1079/SSR2005209>
- Williams PM, Arias I. 1978. Physio-ecological studies of plant species from the arid and semiarid regions of Venezuela. I. The role of endogenous inhibitors in the germination of the seeds of *Cereus griseus* (Haw.) Br.

- and R: (Cactaceae). *Acta Científica Venezolana* **29**: 93-97.
- Willis CG, Baskin CC, Baskin JM, Auld JR, Venable DL, Cavender-Bares J, Donohue K, Rubio-de Casas, RNGWG. 2014. The evolution of seed dormancy: environmental cues, evolutionary hubs, and diversification of the seed plants. *New Phytologist* **203**: 300-309. DOI: <http://dx.doi.org/10.1111/nph.12782>
- Xavier PB, Jasmim JM. 2015. *In vitro* germination and acclimatization of *Hamatocactus setispinus*. *Ornamental Horticulture* **21**: 193-200. DOI: <https://doi.org/10.14295/aohl.v21i2.589>
- Yang XY, Pritchard HW, Nolasco H. 2003. Effects of temperature on seed germination in six species of Mexican Cactaceae. In: Pritchard HW, Probert RJ. eds. *Seed conservation: turning science into practice*. Kew, UK: Royal Botanic Gardens, 575-588. ISBN: 1842460528, 9781842460528
- Yang X, Dong M, Huang Z. 2010. Role of mucilage in the germination of *Artemisia sphaerocephala* (Asteraceae) achenes exposed to osmotic stress and salinity. *Plant Physiology and Biochemistry* **48**: 131-135. DOI: <http://dx.doi.org/10.1016/j.plaphy.2009.12.006>
- Zerpa-Catanho D, Hernández-Pridyballo A, Madrigal-Ortiz V, Zúñiga-Centeno A, Porras-Martínez C, Jiménez VM, Barboza-Barquero L. 2019. Seed germination of pitaya (*Hylocereus* spp.) as affected by seed extraction method, storage, germination conditions, germination assessment approach and water potential. *Journal of Crop Improvement* **33**: 372-394. DOI: <https://doi.org/10.1080/15427528.2019.1604457>
- Zhang H, Irving LJ, McGill C, Matthew C, Zhou D, Kemp P. 2010. The effects of salinity and osmotic stress on barley germination rate: sodium as an osmotic regulator. *Annals of Botany* **106**: 1027-1035. DOI: <http://dx.doi.org/10.1093/aob/mcq204>
- Zimmer K. 1966. UG ber Samengewichte und Samenproduktion einiger Kakteenarten. *Kakteen und andere Sukkulente* **17**: 153-154.
- Zimmer K. 1968. Untersuchungen über den Einfluß der Temperatur auf die Keimung von Kakteen-Saatgut. VII. Über die Keimung einiger argentinischer Arten. *Gartenbauwissenschaft* **33**: 335-344.
- Zimmer K. 1969a. Über die Keimung von Kakteensamen (Schlub). Bedeutung des Lichts und Zusammenfassung. *Kakteen und andere Sukkulente* **20**: 178-180.
- Zimmer K. 1969b. Über die Keimung von Kakteensamen I. Quellung und Keimung bei konstanten Temperaturen. *Kakteen und andere Sukkulente* **20**: 105-107.
- Zimmer K. 1973. Zur Keimung von *Maihuenia poeppigii*. *Kakteen und andere Sukkulente* **24**: 13-14.
- Zimmer K. 1982. Temperatur und keimung einieger kakteenarten. *Kakteen and andere Sukkulente* **33**: 73-75.
- Zimmer K. 1998. Zur Keimung von kakteesaatgut. *Schumannia* **2**: 75-84.
- Zimmer K, Büttner P. 1982. Ersatz des Lichtbedürfnisses bei der Keimung von Kakteensamen durch Gibberellinsäure. *Die Gartenbauwissenschaft* **47**: 97-101.

Associate editor: Arturo de Nova Vázquez

Authors' contributions: DB designed the review, compiled and analyzed literature, made the figures, tables and wrote the first version of the manuscript. JAS, JF and EJ also compiled the literature and contributed to design the manuscript. All authors made multiple revisions and approved the final versions of the manuscript.